

# Host Plant Influence on Activity of *Bacillus thuringiensis* Berliner Against Lepidopterous Pests of Crops<sup>1</sup>

M. Ibrahim Ali, S. Y. Young,<sup>2</sup> and R. C. McNew<sup>3</sup>

Department of Entomology, University of Arkansas, Fayetteville, AR 72701 USA

---

J. Entomol. Sci. 39(3): 311-317 (July 2004)

**Abstract** Mortality of second-instar *Helicoverpa zea* (Boddie), *Spodoptera exigua* (Hübner) and *Pseudoplusia includens* (Walker) fed *Bacillus thuringiensis* Berliner (Bt), Dipel ES®-sprayed leaves of field grown cotton, soybean and tomato were compared in laboratory bioassays. The median lethal concentration (LC<sub>50</sub>) for larvae of all species fed Bt-treated leaf tissue was higher for cotton than for soybean or tomato. The LC<sub>50</sub> for larvae fed Bt-treated soybean and tomato leaves did not differ significantly for any species of insects. When the mean number of days until death was plotted against percent survival at that rate, the mean number of days until death increased with an increase in percent survival. Survival time of the three insect species exposed to Bt increased most rapidly when fed cotton leaves. Results show that foliarly-applied Bt on field-grown cotton leaves is less effective against larvae of these pest species in terms of mortality and speed of kill.

**Key Words** *Bacillus thuringiensis*, host plant, *Pseudoplusia includens*, *Helicoverpa zea*, *Spodoptera exigua*, median lethal concentration

---

*Bacillus thuringiensis* (Bt) is the most commercially successful and widely used biological insecticide in the world (Federici 1999, Lacey et al. 2001). Efficacy is influenced by the host plant for some lepidopterous pests of forest (Appel and Schultz 1994, Moldenke et al. 1994, Farrar et al. 1996, Beveridge and Elek 2001, Kouassi et al. 2001) and field crops (Smith and Hostetter 1982, Meade and Hare 1993, 1994, Faruki and Khan 1996, Luttrell et al. 1998). The mechanistic role of host plant-mediated effects against Bt is related to the quantitative and qualitative variation in allelochemicals in different plant species (Appel and Schultz 1994, Moldenke et al. 1994) and different varieties of some species (Meade and Hare 1993, 1994, Faruki and Khan 1996).

Several examples of host-plant-mediated differences in Bt susceptibility of lepidopterous pests have been reported with deciduous tree species. Efficacy of Bt against *Lymantria dispar* L. was reduced up to five fold on oak, *Quercus* spp., compared to aspen, *Populus tremuloides* Michaux (Appel and Schultz 1994), and showed a >10-fold variation among different host plants (Farrar et al. 1996). Bt-activity was negatively correlated with the concentration of total phenolics, gallotannins, and protein activity in oak leaves (Appel and Schultz 1994). The median lethal concentration

---

<sup>1</sup>Received 15 September 2003; accepted for publication 28 November 2003.

<sup>2</sup>Department of Entomology, 321 AGRI, University of Arkansas, Fayetteville, AR 72701. (email: syoung@uark.edu).

<sup>3</sup>Agricultural Statistics Lab, University of Arkansas, Fayetteville, AR 72701.

(LC<sub>50</sub>) for Bt to the forest tent caterpillar, *Malacosoma disstria* Hübner, was 100-fold greater on aspen than on sugar maple, *Acer saccharum* Marsh (Kouassi et al. 2001). Susceptibility of eucalyptus leaf beetle, *Chrysophtharta bimaculata* (Oliver), to Bt was 104% higher on *Eucalyptus regnans* than *E. nitens* (Beveridge and Elek 2001).

The host plant-mediated alteration of Bt efficacy in field crops is not as well understood. Among field crops, allelochemicals have been shown to significantly reduce growth and survival of lepidopterous larvae on cotton (Chan et al. 1978, Zummo et al. 1984, Hedin et al. 1991) and tomato (Duffey and Isman 1981, Isman and Duffey 1982). Meade and Hare (1994) reported that Bt efficacy against *Spodoptera exigua* (Hübner) and *Trichoplusia ni* (Hübner) was greatest on insects fed resistant lines of celery, *Apium graveolens* var. *rapaceum* (L.). Mortality caused by Bt in *Helicoverpa zea* (Boddie), *Heliothis virescens* (F.), *S. exigua* and *Pseudoplusia includens* (Walker) differed in some instances on cotton and soybean (Luttrell et al. 1998). The survival time of larvae exposed to Bt has been shown to be inversely related to the concentration of Bt (Maczuga and Mierzejewski 1995, Robertson et al. 1996). However, the effect of host plant on survival time of larvae killed by Bt has not been documented. In this study we examined the host plant-mediated variation in susceptibility and time until death of *H. zea*, *S. exigua* and *P. includens* to Bt on cotton, soybean and tomato.

### Materials and Methods

Cotton, *Gossypium hirsutum* L. (cv. DPL 51), soybean, *Glycine max* L. Merrill (cv. DK 5668), and tomato, *Lycopersicon esculentum* L. (cv. Beefsteak) were grown in research plots at the Agricultural Experiment Station, University of Arkansas, Fayetteville, using standard agronomic practices. Neonates of *H. zea* and *P. includens* were obtained from the Insect Rearing Facility in the Entomology Department where they are maintained on artificial diet (Burton 1969). *Spodoptera exigua* eggs were obtained from the USDA Southern Insect Management Research Unit at Stoneville, MS.

Neonates of the test insects were reared to the second stadium on artificial diet in 30-ml clear plastic cups (Solo Cup Company, Urbana, IL). Aqueous suspensions of Dipel® ES (*B. thuringiensis* subsp. *kurstaki*; 17,600 IU potency per mg, Abbott Laboratories, Chemical and Agricultural Product Division, North Chicago, IL) were applied using a spray table on fully expanded terminal leaves of test plants. The spray apparatus was a CO<sub>2</sub>-pressurized boom-type sprayer (Devries Manufacturing, Hollandale, MN) equipped with a Tee Jet 800 2E® hollow-cone nozzle placed 30 cm above the surface of leaves. The apparatus was calibrated to deliver the equivalent of 224 liters/ha. Leaves for the control were sprayed with water. The sprayed leaves were allowed to air dry for approximately 30 min. *Helicoverpa zea*, *P. includens* or *S. exigua* larvae that had been reared to the second stadium on artificial diet were placed individually in 30-ml plastic cups containing one leaf per cup. At least five Bt-rates were tested for each treatment and each treatment was replicated four times. Each group of 20 to 25 larvae in 30-ml cups was placed in a plastic bag containing moist paper towels to assure adequate humidity and held at 28°C. After 2 d, the leaves were searched daily for surviving larvae. Mortality was recorded through pupation. Leaves in the test were replaced every other day with untreated leaves of the same host.

Total mortality in each treatment was analyzed using probit analysis (Proc Probit, SAS Institute 1998). For each insect/host plant combination, the survival time (mean number of days from Bt-exposure until death) at each Bt dose was determined. Only

the larvae that died were considered (Farrar and Ridgeway 1998). The percent survival at each Bt dose was then plotted against the mean time until death (days) of larvae that were killed at that concentration and analyzed to determine their linear relationship using Proc GLM (SAS Institute 1998). Slopes and intercepts for each host plant and insect were compared using Proc GLM Contrast analysis (SAS Institute 1998).

## Results

**Susceptibility of larvae.** *Pseudoplusia includens* was the most susceptible of the 3 species to Bt on all host plants. The  $LC_{50}$  for *S. exigua* was greater than that of *H. zea* on cotton and soybean, but did not differ significantly on tomato. When host plants were compared, the  $LC_{50}$ s for Bt against second instars of all 3 insect species on cotton were at least 6 and 10 fold greater than those on leaves of soybean or tomato plants, respectively. The  $LC_{50}$  did not differ significantly between soybean and tomato for any species tested (Table 1).

**Larval survival time.** Percent larval survival and mean number of days until death for all 3 species fed Bt-treated leaves were positively correlated for all host plants. The slopes of these equations for larval response to Bt-treated cotton were significantly less than for larvae on Bt-treated soybean or tomato, for all 3 insect species. The slope for the larval response to Bt-treated soybean and tomato differed only with *P. includens*, where the slope for soybean was significantly greater (Table 2, Fig. 1).

## Discussion

The  $LC_{50}$  of Bt against three lepidopteran larval pests on leaves of Bt-treated cotton was typically several fold greater than on soybean or tomato leaves. This

**Table 1. The  $LC_{50}$  at pupation of second-instar lepidopteran pests exposed to Dipel ES on leaves of field plants**

| Insect/<br>host plant | No. of<br>larvae | $LC_{50}$ *<br>(mg/ml) | Fiducial limits |        | Intercept | Slope |
|-----------------------|------------------|------------------------|-----------------|--------|-----------|-------|
|                       |                  |                        | Lower           | Upper  |           |       |
| <i>H. zea</i>         |                  |                        |                 |        |           |       |
| Cotton                | 500              | 29.50                  | 23.73           | 38.26  | -2.08     | 1.42  |
| Soybean               | 500              | 2.43                   | 1.96            | 2.93   | -0.87     | 2.26  |
| Tomato                | 480              | 2.84                   | 2.07            | 3.98   | -0.49     | 1.08  |
| <i>P. includens</i>   |                  |                        |                 |        |           |       |
| Cotton                | 500              | 9.25                   | 7.32            | 12.32  | -1.58     | 1.64  |
| Soybean               | 500              | 1.00                   | 0.66            | 1.38   | -0.00     | 1.24  |
| Tomato                | 480              | 0.33                   | 0.00            | 1.02   | 0.60      | 1.24  |
| <i>S. exigua</i>      |                  |                        |                 |        |           |       |
| Cotton                | 550              | 87.68                  | 70.09           | 112.81 | -2.36     | 1.21  |
| Soybean               | 400              | 14.04                  | 5.19            | 41.70  | -1.34     | 1.17  |
| Tomato                | 550              | 4.75                   | 2.06            | 9.89   | -1.08     | 1.49  |

\*  $LC_{50}$ s determined using the Probit Procedure (SAS 1998). Significance was based on overlapping fiducial limits.

**Table 2. Regression equations of percent larval survival and mean number of days until death on Bt-treated leaves of cotton, soybean and tomato**

| Host plant          | Regression equation for larval survival* | Test of significance between slopes of host plants |         | P > F  |
|---------------------|--|--|---------|--------|
| <i>H. zea</i>       |  |  |         |        |
| Cotton              | 18.29 + 11.24 X Mean day                 | Cotton   | Soybean | 0.0475 |
| Soybean             | -80.87 + 35.65 X Mean day                | Soybean  | Tomato  | 0.8401 |
| Tomato              | -74.29 + 38.43 X Mean day                | Tomato   | Cotton  | 0.0334 |
| <i>P. includens</i> |  |  |         |        |
| Cotton              | -31.80 + 26.64 X Mean day                | Cotton   | Soybean | 0.0387 |
| Soybean             | -102.15 + 49.63 X Mean day               | Soybean  | Tomato  | 0.0065 |
| Tomato              | -369.15 + 147.68 X Mean day              | Tomato   | Cotton  | 0.0029 |
| <i>S. exigua</i>    |  |  |         |        |
| Cotton              | 25.45 + 5.11 X Mean day                  | Cotton   | Soybean | 0.0019 |
| Soybean             | -141.54 + 50.85 X Mean day               | Soybean  | Tomato  | 0.1747 |
| Tomato              | -90.16 + 31.43 X Mean day                | Tomato   | Cotton  | 0.0100 |

\* For each insect-host combination, the mean time until death (Mean day) of larvae was calculated for each Bt-dose. Percent survival of larvae at each dose was plotted against the mean days to death to determine their linear relationship using Proc GLM (SAS Institute 1998).

indicates that Bt is less active against larvae on foliage of cotton than on the other host plants. Luttrell et al. (1998) compared these same species of lepidopterous insects on Bt-treated terminals of cotton and soybean, but found little if any difference in susceptibility between host plants. They did find some variability in activity among the different Bt-products for lepidopteran larval pests, *P. includens* and *S. exigua*. The difference in results between the two studies may be due to the different plant part used in that we used fully-developed leaves, while they used terminals. The concentration of certain secondary chemicals such as, tannins, anthocyanin, flavonoids, etc., in fully-developed leaves of cotton plants is greater than in terminal tissues (Zummo et al. 1984, Hedin et al. 1991). Many of these secondary chemicals are bacteriostatic and bactericidal (Smirnoff and Hutchison 1965), and have been shown to reduce the efficacy of Bt-endotoxin against different insects (Maksymiuk 1970, Reichelderfer 1991, Krischik et al. 1988, Appel and Schultz 1994, Navon et al. 1993). The decrease in effectiveness of Bt-treated cotton foliage against larvae that we observed may be related to a greater concentration of these secondary chemicals. Also, protease inhibitors (MacIntosh et al. 1990) may have enhanced the toxicity of Bt against the larvae; these may have been in higher concentration in cotton leaves. Further, the effect of Bt on tomato leaves may have been related to certain allelochemicals (such as chlorogenic acid and polyphenol oxidase) present there. These have been shown to retard larval growth (Elliger et al. 1980, Duffey and Felton 1991, Frelichowski and Juvik 2001) and enhance the efficacy of Bt against *H. zea* larvae (Ludlum et al. 1991).

*Spodoptera exigua* was less affected by Bt than were *H. zea* or *P. includens*. In previous studies on dietary bioassays on artificial diet (Luttrell et al. 1998, 1999), and




Fig. 1. Survival of *H. zea* (A), *P. includens* (B) and *S. exigua* larvae (C) in response to concentration of Bt-applied to foliage of cotton (■), soybean (●) and tomato (▲). For each insect-host combination, the mean time until death (mean day) of larvae was calculated for each concentration. Percent survival of larvae at each dose was plotted against the mean days to determine their linear relationship.

cotton and soybean (Luttrell et al. 1998), the activity of Bt for *S. exigua* was less than for *H. zea* or *P. includens*.

In all tested insect species, percent survival of larvae on Bt-treated leaves was correlated with the mean number of days until larval death. However, this relationship differed somewhat with insect species and with host plants. Larvae fed Bt-treated leaves died faster on tomato and soybean than on cotton. For all three species, the time until death of Bt-treated larvae on field-grown cotton increased more with an increase in percent survival than did that of those on tomato or soybean. This may, however, have been related to secondary chemicals in plants that alter Bt activity or rate of larval development. The phenolics of tomato (Ludlum et al. 1991, Duffey and Felton 1991) or soybean (MacIntosh et al. 1990) enhance efficacy of Bt and may increase the speed of kill. Similarly, leaves of cotton plants contain high concentration of certain phenolics, such as tannins (Chan et al. 1978, Zummo et al. 1984, Hedin et al. 1991) as well as other chemical groups (Chan et al. 1978, Zummo et al. 1984, Hedin et al. 1991) that slow insect development, and may alter Bt activity.

In conclusion, host plant effects must be considered when Bt is applied against pest species on crops. These results suggest that, especially on cotton, Bt may be less active than on some crops and that the speed of killing may differ as well.

### References Cited

- Appel H. M. and J. C. Schultz. 1994.** Oak tannins reduce effectiveness of Thuringide (*Bacillus thuringiensis*) in gypsy moth (Lepidoptera: Lymantriidae). *J. Econ. Entomol.* 87: 1736-1742.
- Beveridge, N. and J. A. Elek. 2001.** Insect and host-tree species influence the effectiveness of a *Bacillus thuringiensis* spp. *tenebrionis*-based insecticide for controlling chrysomelid leaf beetle. *Austr. J. Entomol.* 40: 386-390.
- Burton, R. L. 1969.** Mass rearing of the corn earworm in the laboratory. *USDA, ARS.* 33: 134.
- Chan, B. G., A. C. Weiss, Jr. and M. Lukefahr. 1978.** Condensed tannin, an antibiotic chemical from *Gossypium hirsutum*. *J. Insect Physiol.* 24: 113-118.
- Duffey, S. S. and G. W. Felton. 1991.** Enzymatic antinutritive defenses of the tomato plant against insects, Pp. 166-197. *In* P. A. Hedin, [ed.], *Naturally Occurring Pest Bioregulators.* ACS Symposium Series 44. Am. Chem. Soc., Washington, DC.
- Duffey, S. S. and M. B. Isman. 1981.** Inhibition of insect larval growth by phenolics in glandular trichomes of tomato leaves. *Experientia* 37: 574-576.
- Elliger, C. A., Y. Wong, B. G. Chan and A. C. Weiss, Jr. 1981.** Growth inhibitors in tomato (*Lycopersicon*) to tomato fruitworm (*Heliothis zea*). *J. Chem. Ecol.* 7: 753-758.
- Faruki, S. I. and A. R. Khan. 1996.** The pathogenicity of *Bacillus thuringiensis* var. *kurstaki* on the tropical warehouse moth, *Cadra cautella* (Walker), as influenced by groundnut varieties. *J. Invertebr. Pathol.* 68: 113-117.
- Farrar, Jr. R. R., P. W. A. Martin and R. L. Ridgway. 1996.** Host plant effects on activity of *Bacillus thuringiensis* against gypsy moth (Lepidoptera: Lymantriidae) larvae. *Environ. Entomol.* 25: 1215-1223.
- Farrar, Jr. R. R. and R. Ridgway. 1998.** Quantifying time-mortality relationships for nuclear polyhedrosis viruses when survivors are present. *Environ. Entomol.* 27: 1289-1298.
- Federici, B. A. 1999.** *Bacillus thuringiensis* in biological control, Pp. 575-593. *In* T. S. Bellows and T. W. Fisher [eds.], *Handbook of Biological Control: principle and applications of biological control.* Academic Press, San Diego, CA.
- Frelichowski, Jr. J. E. and J. A. Juvik. 2001.** Sesquiterpene carboxylic acid from a wild tomato species affect larval feeding behavior and survival of *Helicoverpa zea* and *Spodoptera exigua* (Lepidoptera: Noctuidae). *J. Econ. Entomol.* 94: 1294-1259.

- Hedin, P. A., W. L. Parrot and J. E. Jenkins. 1991.** Effects of cotton plant allelochemicals and nutrients on behavior and development of tobacco budworm. *J. Chem. Ecol.* 17: 1107-1121.
- Isman, M. B. and S. S. Duffey. 1982.** Toxicity of tomato phenolics compounds to the fruitworm, *Heliothis zea*. *Entomol. Exp. Appl.* 31: 370-376.
- Kouassi, K. C., F. Lorenzetti, C. Guertin, J. Cabana and Y. Mauffette. 2001.** Variation in the susceptibility of the forest tent caterpillar (Lepidoptera: Lasiocampidae) to *Bacillus thuringiensis* variety *kurstaki* HD-1. Effect of the host plant. *J. Econ. Entomol.* 94: 1135-1141.
- Krischick, V. A., P. Barbosa and C. F. Reichelderfer. 1988.** Three trophic level interactions: Allelochemicals, *Manduca sexta* (L.), and *Bacillus thuringiensis* var. *kurstaki* Berliner. *Environ. Entomol.* 17: 476-482.
- Lacey, L. A., R. Frutos, H. K. Kaya and P. Vail. 2001.** Insect pathogens as biological control agents. Do they have a future? *Biol. Contr.* 21: 130-248.
- Ludlum, C. T., G. W. Felton and S. E. Duffey. 1991.** Plant defenses: Chlorogenic acid and polyphenol oxidase enhance toxicity of *Bacillus thuringiensis* subsp. *kurstaki* to *Heliothis zea*. *J. Chem. Ecol.* 17: 217-236.
- Luttrell, R. G., A. Ali, S. S. Young and K. Knighten. 1998.** Relative activity of commercial formulations of *Bacillus thuringiensis* against selected noctuid larvae (Lepidoptera: Noctuidae). *J. Entomol. Sci.* 33: 365-377.
- Luttrell, R. G., L. Wan and K. Knighten. 1999.** Variation in susceptibility of noctuid (Lepidoptera) larvae attacking cotton and soybean to purified endotoxin proteins and commercial formulations of *Bacillus thuringiensis*. *J. Econ. Entomol.* 92: 21-32.
- MacIntosh, S. C., G. M. Kishore, F. J. Perlak, P. G. Marrone, T. B. Stone, S. R. Sims and R. L. Fuchs. 1990.** Potentiation of *Bacillus thuringiensis* insecticidal activity by serine protease inhibitors. *J. Agric. Food Chem.* 38: 1145-1152.
- Maczuga, S. A. and K. J. Mierzejewski. 1995.** Droplet size and density effects of *Bacillus thuringiensis kurstaki* on gypsy moth (Lepidoptera: Lymantriidae) larvae. *J. Econ. Entomol.* 88: 1376-1379.
- Maksymiuk, B. 1970.** Occurrence and nature of antibacterial substance in plant affecting *Bacillus thuringiensis* and other entomogenous bacteria. *J. Invertebr. Pathol.* 15: 356-371.
- Meade, T. and J. D. Hare. 1993.** Effects of differential host plant consumption by *Spodoptera exigua* (Lepidoptera: Noctuidae) on *Bacillus thuringiensis* efficacy. *Environ. Entomol.* 22: 432-437.
- 1994.** Effects of genetic and environmental host plant variation on the susceptibility of two noctuids to *Bacillus thuringiensis*. *Entomol. Exp. Appl.* 70: 165-178.
- Moldenke, A. F., R. E. Berry, J. C. Miller, J. G. Wernz and X. H. Li. 1994.** Toxicity of *Bacillus thuringiensis* subsp. *kurstaki* to gypsy moth *Lymantria dispar*, fed with alder or douglas fir. *J. Invertebr. Pathol.* 64: 143-145.
- Navon, A., J. D. Hare and B. A. Federici. 1993.** Interactions among *Heliothis virescens* larvae, cotton condensed tannin and the CryIAC  $\delta$ -endotoxin of *Bacillus thuringiensis*. *J. Chem. Ecol.* 19: 2485-2499.
- Reichelderfer, C. F. 1991.** Interactions among allelochemicals, some lepidoptera, and *Bacillus thuringiensis* Berliner, Pp. 507-524. *In* P. Barbosa, V. E. Krischik and Clive G. Jones [eds.], *Microbial Mediation of Plant-Herbivore Interaction*. John Wiley & Sons, Inc. NY.
- Robertson, J. L., H. K. Preisler, S. S. Ng, L. A. Hickie, A. Berdeja and W. D. Gelernter. 1996.** Comparative effect of temperature and time on activity of Dipel 2x and MVP preparations of *Bacillus thuringiensis* subsp. *kurstaki* on diamondback moth (Lepidoptera: Plutellidae). *J. Econ. Entomol.* 59: 1084-1087.
- SAS Institute Inc. 1998.** SAS/STAT User's Guide, Release 6.03 Edition. Cary, NC.
- Smith, D. B. and D. L. Hostetter. 1982.** Laboratory and field evaluations of pathogen-adjuvants. *J. Econ. Entomol.* 75: 472-476.
- Smirnov, W. A. and P. M. Hutchison. 1965.** Bacteriostatic and bacteriocidal effects of extracts of foliage from various plant species on *Bacillus thuringiensis* var. *thuringiensis* Berliner. *J. Invertebr. Pathol.* 7: 273-280.
- Zummo, G. R., J. C. Segers and J. H. Benedict. 1984.** Seasonal phenology of allelochemicals in cotton and resistance to bollworm (Lepidoptera: Noctuidae). *Environ. Entomol.* 13: 1287-1290.