

Influence of Trap Color on Collection of the Recently-Introduced Bean Plataspid, *Megacopta cribraria* (Hemiptera: Plataspidae)¹

Scott Horn² and James L. Hanula

USDA Forest Service, Southern Research Station, Forestry Sciences Laboratory, 320 Green Street, Athens, Georgia 30602-2044 USA

J. Entomol. Sci. 46(1): 85-87 (January 2011)

Large numbers of the exotic bean plataspid, *Megacopta cribraria* (F.) (Hemiptera: Plataspidae), were first collected from several northeast Georgia counties beginning in October 2009 (Suiter and Ames 2009, Statewide Pest Alert). How this insect arrived in the United States and where it came from is still not known. The native range of *M. cribraria* is reported to be throughout Asia and the Indian Subcontinent (Hosokawa et al. 2007, Proc. R. Soc. B. 274: 1979 - 1984; Srinivasaperumal et al. 1992, Proc. Indian Natn. Sci. Acad. 6: 333 - 340; Hua 2000, List of Chinese Insects. Vol. 1, Zhong-shan Univ. Press, Guangzhou. 251 pp.). It is similar to other Plataspidae in having a somewhat unusual symbiotic relationship with its gut bacteria. Before laying eggs, females deposit particles containing the symbiont which are then eaten by newly-hatched nymphs under natural conditions. Nymphs experimentally deprived of access to the symbiont exhibited slower growth, smaller body sizes, and higher mortality. Thus, presence of the obligate symbiont was predictive of the insect's status as a pest (Hosokawa et al. 2007). Preliminary testing of *M. cribraria* in Georgia showed that all individuals evaluated contained the symbiont (Jenkins et al. 2010, J. Entomol. Sci. 45: 62 - 63).

The plant hosts of the Plataspidae are predominantly legumes (i.e., bean family) (Schaefer and Panizzi 2000, Heteroptera of Economic Importance. CRC Press, Boca Raton, FL. 828 pp.), such as kudzu (*Pueraria montana* (Lour.) Merr.), lablab bean (*Lablab purpureus* L.), and soybeans (*Glycine max* (L.) Merr.). Although the occurrence of a new insect feeding on kudzu was welcomed in Georgia, the possibility of an insect damaging any portion of the \$12.5 billion U.S. soybean crop is not. Reports from China suggest *M. cribraria* can cause serious damage to soybean production (Zhixing et al. 1996, Plant Protect. 3: 7 - 9; Guangnan et al. 2006, Acta Agro. Sinica. 32: 491 - 496.). In addition to soybeans, other legume crops have been negatively affected by this pest such as *lablab* beans (Thippeswamy and Rajagopal 1998, Karnataka J. Ag. Sci. 11: 941 - 946) and *Phaseolus* beans (Easton and Pun 1997, Proc. Entomol. Soc. Wash. 99: 574 - 582). Furthermore, many important forest plant species in the bean family (i.e., redbud, black locust) can be found in the southeastern U.S., and how they might be impacted by this pest remains unknown.

¹Received 23 July 2010; accepted 7 October 2010.

²Corresponding author (shorn01@fs.fed.us).



Fig. 1. Cross-vane trap used to collect adult *M. cribraria*.

Establishing baseline information on abundance and seasonal activity is an important first step in evaluating and studying a new insect in a novel environment. Because *M. cribraria* (and the family Plataspididae) is new to North America, no studies exist on how to best collect and monitor the species. The effect of trap color on collection performance has been studied extensively for many groups of insects such as lepidopterans (Roubos and Liburd 2008, J. Agric. Urban Entomol. 25: 99 - 109), coleopterans (Al-Saoud et al. 2010, J. Entomol. 7: 54 - 59), and hymenopterans (Campbell and Hanula 2007, J. Insect Cons. 11: 399 - 408; Stephen and Rao 2005, J. Kansas Entomol. Soc. 78: 373 - 380). Species often show strong attraction to certain colors which permits the development of selective monitoring traps for them (Kirk 1984, Ecol. Entomol. 9: 35 - 41). Of particular interest here are the hemipterans, adults of which are typically monitored in agricultural environments using yellow traps. For example, false chinch bug (*Nysius raphanus* Howard) in Colorado (Cranshaw 2006, Phytoparasitica 34: 197 - 203), plant bugs (*Lygus* spp.) in Washington (Landis and Fox 1972, Environ. Entomol. 1: 464 - 465), and psyllids in West Virginia (Cooper et al. 2010, Can. Entomol. 142: 188 - 191) were all highly attracted to yellow traps.




Fig. 2. Mean number (\pm SE) of adult *M. cribraria* captured in different colored cross-vane traps. Bars with the same letter are not significantly different ($P < 0.05$) according to the REGWQ.

Our objective was to determine the effect of trap color on catches of adult *M. cribraria*. The study was conducted in a kudzu field approx. 30 ha in size in Athens, GA. Early-season evaluations of the field showed large numbers of over-wintered adults with females beginning to lay eggs. On 10 May 2010, we established 4 rows each containing a white, yellow, purple, red, and black cross-vane trap. Rows were spaced approx. 50 m apart and traps were 5 m apart within rows. We used plexiglass cross-vane traps (Fig. 1) identical to those of Ulyshen and Hanula (2007, Am. Midl. Nat. 158: 260 - 278) but our buckets were gray instead of white and the plexiglass panels were spray painted one of the 5 colors. Traps were operated continuously from 10 May to 5 July 2010 and checked weekly. Collection buckets were filled with a soapy-water solution and all adult *M. cribraria* collected in them were counted in the field. Data were analyzed using a two-way analysis of variance (ANOVA) (SAS Institute 1982). Means separation was achieved using the Ryan-Einot-Gabriel-Welsch (REGWQ) multiple range test. We used the $\log_{10}(x + 1)$ transformation to reduce heteroscedasticity (Sokal and Rohlf 1981).

A total of 4,057 *M. cribraria* were collected, and color significantly affected trap catches ($F = 47.00$; $df = 4, 12$; $P < 0.0001$). White traps captured over twice the number of *M. cribraria* as yellow traps and both colors captured significantly more than black, purple, or red (Fig. 2). Many monitoring programs use yellow traps for collection of phytophagous pests. Although yellow traps were effective for capturing the bean plataspid, white traps were much more effective capturing 53% of all *M. cribraria* collected in our study. Therefore, we suggest utilizing white traps to increase the likelihood of early detection and for monitoring relative abundance of this insect.

Acknowledgments

The authors thank Mike Cody for help with field work and Yanzhuo Zhang for reviewing the manuscript.

AUTHOR GUIDELINES

The *Journal of Entomological Science* (JES), published quarterly by the Georgia Entomological Society, invites submission of manuscripts reporting original research with insects and related arthropods or literature reviews offering foundations to innovative directions in entomological research.

Submission and Review. Three (3) copies of all manuscripts to be submitted for consideration for publication in JES should be sent to **Wayne Gardner, Editor, *Journal of Entomological Science*, 1109 Experiment Street, UGA Griffin Campus, Griffin, GA 30223-1797 USA.** Manuscripts judged as suitable for JES will be peer reviewed by scientists qualified to assess the scientific merit of the content. The Editor has the final decision in the acceptance, rejection, or revision of submitted/reviewed manuscripts.

Publication Charges. Authors will be charged US\$55 per published page. Pages containing figures will be subject to a US\$20 surcharge. Additional costs incurred in publishing color figures or photos must be passed onto the author. These costs vary with several factors, and every effort will be made to reduce these costs. Page charges will be reduced US\$15 per page for those articles with senior authors who are members in good standing with the Georgia Entomological Society. All fees are subject to change without notice.

Style and Format. Full guidelines for manuscripts can be found at www.ent.uga.edu/ges. Lines within text and tables of manuscripts should be double-spaced. A **title page** must contain the name, address, telephone number and email address of the corresponding author in the upper right. The running head, title, author(s), institutional affiliation where the research was conducted, and any footnotes should be left justified on the remainder of the title page. An **Abstract** followed by **key words** must follow on a separate page. The **text** should be divided into an introduction (no heading), Materials and Methods, Results and Discussion (together or separate), and an Acknowledgment(s) (optional). The **References Cited** must begin on a new page, and the **figure captions** and individual **tables** and **figures** must be placed on separate pages at the end of the text. All headings, subheadings, and table and figure captions should be in boldface font.

Figure Preparation. For review purposes, figures may be black and white or color with each figure on a separate page. However, figures in final revised typescripts must be in one of the following formats: TIFF, EPS, WMF, JPEG, or PowerPoint. (GIF formats are unacceptable as they produce poor quality images.) Although figures of any size can be submitted, those that fit the width of the printed page (118 mm) expedite the publication process. In choosing font size of labels and legends, insure that these are sufficiently large so that in reducing figures to fit the printed page (118 × 188 mm), lettering will not be difficult to read. Photographs should be submitted in TIFF format. Black and white photos should be in grayscale, not color. The CMYK color mode—not RGB—must be used for color photos.

Notes. JES also publishes Notes of original research and observations that may not be sufficiently replicated for acceptance as Scientific Papers but have merit in terms of sharing with the scientific community. These are designed to be short and rapid communications with none to few tables or figures. Notes do not contain an abstract or separate text sections (e.g., introduction, materials and methods, results and discussion). References are cited within the text by placement of authors' last names, date, and journal or book citation within parentheses.