

Global Gene Expression in Cotton Fed Upon by *Aphis gossypii* and *Acyrtosiphon gossypii* (Hemiptera: Aphididae)¹

Quancheng Zhang², Yudong Zhang², and Jungang Wang³

College of Agriculture, Shihezi University, Shihezi 832003, China

J. Entomol. Sci. 58(1): 47–68 (January 2023)

DOI: 10.18474/JES22-07

Abstract *Aphis gossypii* Glover and *Acyrtosiphon gossypii* Mordvilko (Hemiptera: Aphididae) are key pests of cotton, *Gossypium hirsutum* L., known to induce cotton host plant defense responses. Deep RNA sequencing of the cotton transcriptome followed by differential expression analyses were performed to clarify the molecular mechanisms of cotton defense in response to feeding by these aphid pests. We found 6,565 genes were differentially expressed in cotton in response to feeding by *Ac. gossypii* and 823 genes that were differentially expressed in response to feeding by *A. gossypii*, while 2,379 genes were differentially expressed in response to simultaneous feeding by both species. Pathway enrichment analysis showed that the differentially expressed genes associated with *Ac. gossypii* feeding were enriched for metabolic pathways, porphyrin and chlorophyll metabolism, biosynthesis of secondary metabolites, biosynthesis of carotenoids, and the pentose phosphate pathway. The enriched pathways in cotton fed on by *A. gossypii* were thiamine metabolism, glutathione metabolism, plant–pathogen interaction, and sesquiterpene and triterpenoid biosynthesis. The differentially expressed genes in cotton induced by simultaneous feeding of both species were primarily related to circadian rhythm regulation, photosynthesis, porphyrin and chlorophyll metabolism, galactose metabolism, and flavonoid biosynthesis.

Key Words *Aphis gossypii*, *Acyrtosiphon gossypii*, cotton, transcriptome, gene expression

Cotton, *Gossypium hirsutum* L., is an economically important crop in global agricultural and textile industries. Aphids (Hemiptera: Aphididae) have become important pests of cotton production worldwide and are now considered the dominant pest species of cotton-growing areas in China (Lu and Liang 2016, Lu et al. 2020).

Although insects with piercing-sucking mouthparts cause less mechanical damage to host plants than insects with chewing mouthparts, the damage cycle is longer, resulting in plants exhibiting mild but persistent defense responses to the attack (Moran and Thompson 2001, Thompson and Goggin 2006). Plants possess specialized structures and substances, for example, wax, hairs, spines, glands, and different degrees of ossification or silicification of some tissues on the plant surface, as defense mechanisms (Zhang et al. 2013c). While these specialized tissues and

¹Received 7 February 2022; accepted for publication 2 April 2022.

²Co–first authors.

³Corresponding author (email: wangjungang98@163.com).

structures help to resist pests, they cannot completely defend the host plant from attack by phytophagous insects.

Insects with piercing-sucking mouthparts can penetrate the plant epidermis to imbibe plant fluids from the phloem and xylem. This may lead to depolarization of the lipid membrane or a disturbance in the transmembrane ion flow in plant cells, causing a change in transmembrane potential across the cell membrane and a change in signal transduction: for example, the calcium ion (Ca^{2+}) influx (Bricchi et al. 2012, Luo et al. 2017, Vincent et al. 2017, Yan et al. 2018). Plants also regulate the activity of key proteases to degrade and eliminate reactive oxygen chemical species, including phenolic and quinone compounds (Chen et al. 2011, Liu and Lan 2009, Luo et al. 2008, Tjallingii 2006, Wang et al. 2011, Wu et al. 2015). This defense mechanism likely evolved while resisting the chemicals injected during feeding of insects with piercing-sucking mouthparts (Boyko et al. 2006, Voelckel et al. 2004, Zhou et al. 2009).

Aphis gossypii Glover and *Acyrtosiphon gossypii* Mordvilko (Hemiptera: Aphididae) are common pests of cotton in China and worldwide. Their feeding may cause leaf curl, stunting of plant growth, and slowing of plant development (Jacobson and Croft 1998). Honeydew produced while feeding may serve as a nutritional substrate for molds that can interfere with light absorption and photosynthetic activity (Hullé et al. 2020). Cotton plants also exhibit a series of physiological and metabolic reactions with stress associated with *A. gossypii* and *Ac. gossypii* feeding. These include increased activity of the antioxidative enzymes catalase (CAT), peroxidase (POD), polyphenol oxidase (PPO), lipoxygenase (LOX), and other defense enzymes (Chen et al. 2015; Li et al. 1998a, b).

The alteration in the level of soluble sugars, free proline, and other nutrients due to aphid attack initiates immune defense mechanisms against aphid feeding (Li et al. 2008, Patima et al. 2018). We have previously shown that *A. gossypii* and *Ac. gossypii* feeding can cause various defense responses in cotton. For example, *A. gossypii* was found to cause changes in chlorophyll, soluble protein, proline, malondialdehyde content, and antioxidant enzyme activity in cotton at both the boll and bud stages and, with the extension of stress time, cotton defense ability was enhanced (Deng et al. 2013, Yan et al. 2013). Feeding by *Ac. gossypii* altered the level of soluble sugar, soluble protein, chlorophyll, carotenoids, malonaldehyde, and the activity of POD in cotton; nutrient metabolism and cell permeability also were altered. At the same time, the activity of related defense enzymes was induced (Zhang et al. 2020). However, the specific gene expression changes that mediate cotton defense responses to *A. gossypii* and *Ac. gossypii* attacks remain poorly understood. The aim of this study was, therefore, to use transcriptome sequencing to investigate the differential expression of genes related to biological processes, cell components, and molecular functions in cotton following feeding by *A. gossypii* and *Ac. gossypii*.

Materials and Methods

Experimental treatments. Cotton (New Upland Early Maturity 44 variety) seeds were soaked for 1 h at 55°C, allowed to germinate at room temperature for 24 h, and then planted in vermiculite in plastic basins (12-cm height, 10-cm diameter), which

were maintained in environmentally controlled incubators on a 16:8-h photoregime at 24°C from midnight to 8:00 a.m. and 27°C from 8:00 a.m. to 11:59 p.m. Tests with aphids were initiated when cotton seedlings had grown to two true leaves.

Aphis gossypii and *Ac. gossypii* aphids used in the study were from colonies that had been subcultured on cotton seedlings for more than 30 generations. Aphids from these colonies were transferred individually to cotton seedlings using a fine brush. Nine seedlings were infested with 16 *Ac. gossypii* per plant; nine were infested with 16 *A. gossypii* per plant; nine were infested with eight *Ac. gossypii* and eight *A. gossypii* per plant; and nine were not infested and served as a check. Once aphids were transferred to the plants, whole plants were covered and placed in an incubator with controlled lighting.

After 3 d, the aphids on the cotton seedlings were removed, and the new plant growth at the top of each plant was excised with scissors and placed individually in 1.5-ml sterile centrifuge tubes. These were immediately placed in liquid nitrogen and transferred to -80°C. Each of the four treatments was replicated three times.

RNA extraction, sequencing, and data analysis. Sample RNA extraction, quality detection, transcriptome sequencing, and statistical analyses were commissioned and performed by Beijing Nuohe Zhiyuan Technology Co., Ltd. (Beijing, China). Briefly, total RNA was extracted from the cotton samples using TRIzol (Tiangen Biotech [Beijing] Co., Ltd., Beijing, China) according to manufacturer's instructions. RNA purity was checked using the NanoPhotometer® spectrophotometer (Implen, Corston, United Kingdom). RNA concentration was measured using the Qubit® RNA Assay Kit in Qubit® 2.0 Fluorometer (Life Technologies, Carlsbad, CA), and RNA integrity was assessed with the RNA Nano 6000 Assay Kit of the Bioanalyzer 2100 system (Agilent Technologies, Santa Clara, CA). All samples had an RNA integrity number (RIN) above 6.7. RNA sequencing libraries were prepared with NEBNext® Ultra™ RNA Library Prep Kit for Illumina® (Illumina, Inc., San Diego, CA) and sequenced on an Illumina HiSeq™ 2500 platform at an average depth of ~66 million reads per sample. Raw sequencing reads were quality assessed with FastQ. To pass the initial quality control check, the average Phred score of each base position across all reads had to be at least 30. Reads were further processed by cutting individual low-quality bases and removing adapter and other Illumina-specific sequences with ng-qc using default parameters. HISAT2-2.0.4 was then used to map the trimmed reads to the cotton AD1_ZJU_v2.1 reference genome (Kim et al. 2015, Mortazavi et al. 2008). To quantify gene expression levels, mapped reads were summarized at the gene level using HTSeq version 0.6.0 (Anders 2010). Differential expression analyses were performed with DESeq2 R package (version 1.10.1), and gene ontology (GO) enrichment analyses were conducted using the clusterProfiler R package (Anders and Huber 2012, Wang et al. 2010). The significance threshold used was the adjusted *P* value of 0.05 and absolute fold change of 2 for the differential expression analysis and adjusted *P* value less than 0.05 for GO enrichment analysis (Robinson et al. 2010, Young et al. 2010). We used clusterProfiler R package to test the statistical enrichment of differential expression genes in the Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways (Kanehisa et al. 2008).

Table 1. Results of RNA quality testing on cotton leaves fed on by aphids.

Sample*	Concentration (ng/μl)	Total (μg)	Optical Density 260/280 Value	Optical Density 260/230 Value	RIN Value	Result
Control	452	14.464	1.78	1.78	7.3	Qualification
Control	418	13.376	1.73	1.38	6.9	Qualification
Control	306	9.792	1.76	1.78	7.5	Qualification
Acy1	314	10.048	1.73	1.67	7.1	Qualification
Acy2	334	10.688	1.70	1.61	7.0	Qualification
Acy3	264	8.448	1.76	1.78	6.7	Qualification
A1	306	9.792	1.78	0.68	7.0	Qualification
A2	356	11.392	1.73	1.60	6.9	Qualification
A3	414	13.248	1.80	1.92	6.8	Qualification
AcyA1	822	26.304	1.86	1.44	7.3	Qualification
AcyA2	474	15.168	1.80	1.99	7.5	Qualification
AcyA3	252	8.064	1.64	1.43	7.4	Qualification

* A, *Aphis gossypii*; Acy, *Acyrtosiphon gossypii*; AcyA, *Acyrtosiphon gossypii* and *Aphis gossypii*; CK, XXX; OD, optical density; RIN, RNA integrity number.

Results

RNA quality and sequencing data results. All indicators of RNA quality (260/280 and 260/230 absorbance ratios and RIN values) were above accepted quality thresholds for all samples, and the extracted RNA could therefore be used for transcriptome sequencing (Table 1). Sequencing data statistics showed that for each sample the sequencing data error rate was $\leq 0.03\%$; $>97\%$ of sequences had a Phred score of at least Q20, $>92\%$ of sequences had a Phred score of at least Q30, and the guanine and cytosine (GC) content was stable around 44%, indicating that the sequencing data had sufficient quality to be used for subsequent analyses (Table 2). Furthermore, the average sequencing depth was >60 million reads per sample, and the filtered sequencing reads had a high alignment with the reference genome, suggesting an appropriate sequencing depth for differential expression analyses (Table 3).

Gene expression analysis. According to the results of gene expression in cotton fed upon by *A. gossypii* and *Ac. gossypii* (Fig. 1), the fragments per kilo base of exon per million reads (FPKM) values of gene expression were divided into five levels. Within the range of FPKM values of 0–15, the amount of gene expression in cotton had no significant relationship with the feeding of *A. gossypii* and *Ac. gossypii*. In the range of FPKM values of 15–60, *A. gossypii* feeding had no significant effect on cotton gene expression compared to the control group, while

Table 2. Statistical results for cotton transcriptome sequencing data after aphid feeding.

Sample*	Raw Reads	Clean Reads	Clean Bases	Error Rate (%)	Phred Score (%)		Guanine-Cytosine Content (%)
					Q20	Q30	
Control	57608420	56981220	8.55G	0.03	97.54	92.80	44.44
Control	52977718	52398266	7.68G	0.03	97.62	92.92	44.47
Control	56398218	55843658	8.38G	0.03	97.60	92.89	44.30
Acy1	67742536	67010136	10.05G	0.03	97.58	93.00	44.81
Acy2	75035796	74217572	11.13G	0.03	98.08	94.05	44.57
Acy3	81274336	80530890	12.08G	0.02	98.13	94.17	44.55
A1	69087066	68516836	10.28G	0.02	98.24	94.42	44.34
A2	79433746	78799342	11.82G	0.03	98.10	94.09	44.63
A3	83596158	82856886	12.43G	0.02	98.15	94.24	44.72
AcyA1	60617246	59897002	8.98G	0.03	98.04	94.05	44.91
AcyA2	60026858	59380680	8.91G	0.02	98.23	94.49	44.62
AcyA3	56710670	55949342	8.39G	0.03	97.46	92.61	44.44

* A, *Aphis gossypii*; Acy, *Acyrtosiphon gossypii*; AcyA, *Acyrtosiphon gossypii* and *Aphis gossypii*; CK, XXX; GC, XXX; G, XXX.

feeding by *Ac. gossypii* alone and in combination with *A. gossypii* resulted in significantly reduced cotton gene expression. In the range of FPKM value >60, the amount of gene expression in cotton fed on by *Ac. gossypii* was significantly lower compared to the other groups (Fig. 1).

Differential gene expression analysis. An overview of the differentially expressed genes in cotton fed upon by *A. gossypii* and *Ac. gossypii* is shown in Table 4. A total of 6,565 genes (3,310 genes upregulated and 3,255 genes downregulated) were differentially expressed between cotton fed on by *Ac. gossypii* and cotton free of aphids (Fig. 2A); 823 genes (470 upregulated and 353 downregulated) were differentially expressed between cotton fed upon by *A. gossypii* and cotton free of aphids (Fig. 2B); and 2,379 genes (1,003 upregulated and 1,376 downregulated) were differentially expressed between cotton fed upon by *A. gossypii* and cotton fed upon by *Ac. gossypii* (Fig. 2C). The upregulated genes in cotton fed upon by *Ac. gossypii* were mainly concentrated in the photosynthetic metabolism pathway, biosynthesis of secondary metabolites, and the pentose phosphate metabolism pathway. The upregulated genes in cotton fed upon by *A. gossypii* were mainly concentrated in amino acid metabolism, plant-pathogen interaction, and terpene biosynthesis. When mixed populations of *A. gossypii* and *Ac. gossypii* fed on the cotton, the upregulated genes were mainly concentrated in circadian rhythm regulation, photosynthesis, and galactose metabolism pathways (Fig. 3). Among these differentially expressed genes, 280

Table 3. Analysis of reference sequences of cotton transcriptome sequencing data for samples fed on by aphids.

Sample*	Total Reads	Total Mapped (%)	Multiple Mapped (%)	Uniquely Mapped (%)	Splice Read (%)
CK	56,981,220	55,333,069 (97.11)	2,332,815 (4.09)	53,000,254 (93.01)	20,512,520 (36)
CK	52,398,266	50,987,846 (97.31)	2,120,254 (4.05)	48,867,592 (93.26)	18,861,344 (36)
CK	55,843,658	54,183,366 (97.03)	2,410,084 (4.32)	51,773,282 (92.71)	20,335,462 (36.41)
Acy1	67,010,136	65,075,646 (97.11)	2,876,427 (4.29)	62,199,219 (92.82)	23,956,281 (35.75)
Acy2	74,217,572	72,408,668 (97.56)	2,943,500 (3.97)	69,465,168 (93.6)	26,356,169 (35.51)
Acy3	80,530,890	78,732,584 (97.77)	3,122,866 (3.88)	75,609,718 (93.89)	28,310,160 (35.15)
A1	68,516,836	66,765,122 (97.44)	2,694,788 (3.93)	64,070,334 (93.51)	24,595,724 (35.9)
A2	7,8799,342	76,669,137 (97.3)	3,179,895 (4.04)	73,489,242 (93.26)	28,591,583 (36.28)
A3	82,856,886	80,129,634 (96.71)	3,870,914 (4.67)	76,258,720 (92.04)	29,500,911 (35.6)
AcyA1	59,897,002	57,010,099 (95.18)	2,994,953 (5)	54,015,146 (90.18)	19,613,832 (32.75)
AcyA2	59,380,680	58,017,172 (97.7)	2,436,567 (4.1)	55,580,605 (93.6)	21,509,592 (36.22)
AcyA3	55,949,342	54,413,783 (97.26)	2,406,065 (4.3)	52,007,718 (92.96)	19,877,962 (35.53)

* A, *Aphis gossypii*; Acy, *Acyrtosiphon gossypii*; AcyA, *Acyrtosiphon gossypii* and *Aphis gossypii*; CK, XXX.




Fig. 1. Cotton gene expression profile after feeding by *Acyrtosiphon gossypii* (Acy) and/or *Aphis gossypii* (A). FPKM, reads per kilobase million mapped reads. Different letters on the column indicate significant differences between treatment groups ($P < 0.05$). There were differences in each treatment when FPKM interval was 15–60 and >60 ($P < 0.05$), and there was no significant difference when FPKM interval was 0–1, 1–3, and 3–15 ($P > 0.05$).

were common in the first two comparisons, and 96 genes were shared across all three comparisons (Fig. 4).

GO enrichment analysis of differentially expressed genes. *Aphis gossypii* feeding on cotton resulted in regulation of biological processes related to synthesis and metabolism of tetraterpenoid, carotenoid, and methionine, and molecular functionals related to flavin adenine dinucleotide (FAD) binding and the activity of 5-

Table 4. Number of genes in cotton induced by aphids.

Comparison Combination*	Differential Gene	Upregulated Genes	Downregulated Genes
Acy versus CK	6,565	3,310	3,255
A versus CK	823	470	353
A versus Acy	7,256	3,695	3,561
AcyA versus CK	2,379	1,003	1,376
AcyA versus Acy	797	323	474
AcyA versus A	2,474	890	1,584

* A, *Aphis gossypii*; Acy, *Acyrtosiphon gossypii*; AcyA, *Acyrtosiphon gossypii* and *Aphis gossypii*; CK, XXX.




Fig. 2. Volcano plot of differentially expressed genes. (A) Volcano plot of differentially expressed genes after feeding by *Acyrtosiphon gossypii* (Acy). (B) Volcano plot of differentially expressed genes after feeding by *Aphis gossypii* (A). (C) Volcano plot of differentially expressed genes after feeding by both of *Acyrtosiphon gossypii* and *Aphis gossypii* (AcyA). Red dots indicate upregulated genes, green dots indicate downregulated genes ($P > 0.05$), blue dots indicate nonsignificant genes ($P < 0.05$).




Fig. 3. Cluster analysis of differentially expressed genes in cotton after feeding by *Acyrtosiphon gossypii* (Acy) and/or *Aphis gossypii* (A). The $\log_{10}$ (fragments per kilo base of exon per million reads [FPKM] + 1) value was normalized and transformed (scale number) and clustered. Red represented the high-expression gene, and blue represented the low-expression gene. Color from red to blue, representing $\log_{10}(\text{FPKM} + 1)$ in descending order.

methyltetrahydropteroyltriglutamate-homocysteine S-methyltransferase and oxidoreductase, among others (Fig. 5A). *Acyrtosiphon gossypii* feeding on cotton had a modulatory effect on genes involved in photosynthesis, steroid biosynthesis, lipid biosynthesis, and other biological processes, as well as cellular components (e.g., thylakoid, photosystem II oxygen complex, oxidoreductase complex, and cell




Fig. 4. Venn diagrams showing the number of differentially expressed genes in cotton after feeding by *Acyrtosiphon gossypii* (Acy) and/or *Aphis gossypii* (A). The sum of the numbers in each large circle represents the total number of differential genes in the comparison combination, and the overlapping part of the circle represents the common differential genes between the combinations.

membrane components in the photosynthetic system). The expression of genes related to molecular functions, such as oxidoreductase activity and steroid dehydrogenase activity, were also affected (Fig. 5B). Feeding by mixed populations of *A. gossypii* and *Ac. gossypii* had a significant effect on biological processes related to phosphorus signal transduction in cotton (Fig. 5C).

KEGG enrichment analysis of differentially expressed genes. In the process of cotton differential gene enrichment, porphyrin and chlorophyll metabolism were the most enriched after feeding by *Ac. gossypii*, with 41 genes in the pathway. The second was monoterpenoid biosynthesis, in which six genes were involved in this pathway, and the least enriched were metabolic pathways (Fig. 6A). Thiamine metabolism was the most enriched in cotton after feeding by *A. gossypii*, with five genes in the pathway. The second was diterpenoid biosynthesis, which had four genes. Biosynthesis of secondary metabolites was the least enriched (Fig. 6B). Plant circadian rhythm was the most enriched in cotton after feeding by both *Ac. gossypii* and *A. gossypii*, with 26 genes in the pathway. The second was thiamine metabolism, which had five genes. The least enrichment was in metabolic pathways (Fig. 6C).

We found that *Ac. gossypii* feeding on cotton resulted in regulation of 1,221 annotated genes (Table 5), and five KEGG pathways were significantly enriched, including metabolic pathways, porphyrin and chlorophyll metabolism, biosynthesis of secondary metabolites, biosynthesis of carotenoids, and the pentose phosphate pathway (Table 6). The numbers of differentially expressed genes annotated in metabolic pathways and biosynthesis of secondary metabolites were 723 and 403, respectively. Thirty-eight annotated genes were differentially expressed after *A. gossypii* feeding (Table 5), and four enriched pathways were identified, including




Fig. 5. Gene ontology (GO) enrichment column diagram. (A) GO enrichment analysis of cotton differentially expressed genes after feeding by *Acyrtosiphon gossypii* (Acy); (B) GO enrichment analysis of cotton differentially expressed genes after feeding by *Aphis gossypii* (A); (C) GO enrichment analysis of cotton differentially expressed genes after feeding by both of *Acyrtosiphon gossypii* and *Aphis gossypii* (AcyA). Different colors to distinguish biological processes, cellular components, and molecular functions. GOrms plot with “*” for significant enrichment.




Fig. 6. Enrichment scatter plot of differential gene KEGG. (A) The differential gene KEGG in cotton was enriched feeding by *Acyrtosiphon gossypii* (Acy). (B) The differential gene KEGG in cotton was enriched feeding by *Aphis gossypii* (A). (C) The differential gene KEGG in cotton was enriched feeding by both of *Acyrtosiphon gossypii* and *Aphis*

Table 5. KEGG functional annotation of differentially expressed genes in cotton induced by aphids.

Comparison Combination*	No. of Annotated Genes	No. of Annotated Differential Genes	No. of Pathways Annotated	No. of Pathways Enriched
Acy versus CK	3,091	1,224	83	5
A versus CK	362	38	114	4
A versus Acy	3,307	1,206	114	4
AcyA versus CK	1,129	355	101	6
AcyA versus Acy	343	44	79	4
AcyA versus A	1,116	75	104	4

* A, *Aphis gossypii*; Acy, *Acyrtosiphon gossypii*; AcyA, *Acyrtosiphon gossypii* and *Aphis gossypii*; CK, XXX.

thiamine metabolism, glutathione metabolism, plant–pathogen interaction, and sesquiterpene and triterpenoid biosynthesis (Table 6). Among these, the plant–pathogen interaction pathway included the largest number of annotated genes (16). A total of 355 annotated genes were differentially expressed after feeding by mixed populations of *A. gossypii* and *Ac. gossypii* (Table 5); six KEGG pathways were significantly enriched, including circadian rhythm regulation, photosynthesis, porphyrin and chlorophyll metabolism, galactose metabolism, and flavonoid biosynthesis (Table 6).

Discussion

Plants attacked and fed upon by insects with piercing-sucking mouthparts activate related resistance genes (Park et al. 2005), induce an emergency response to the injury at the feeding site, activate the whole-plant defense system to reduce plant damage, and prepare for a rapid defense response to future disturbances (Martinez-Medina et al. 2016). Indeed, host plant defense regulatory mechanisms may differ with insect species, developmental stage, or feeding mechanism (chewing versus piercing-sucking mouthparts), as well as host plant species and characteristics. Yet, all play an important role in host plant defense against pests.

***gossypii* (AcyA).** The size of the point indicates the number of differentially expressed genes in this pathway, and the color of the point corresponds to different Q value ranges. The greater the Rich factor, the greater the degree of enrichment. Q value is the *P* value after multiple-hypothesis testing correction. The range of the Q value is [0,1]: the closer to zero, the more obvious enrichment.

Table 6. Pathway enrichment of differentially expressed genes in cotton induced by aphids.

Comparison Combination*	Term	Input No.	Background No.	Corrected P Value
Acy versus CK	Metabolic pathways	723	1,861	0.0000434
	Porphyrin and chlorophyll metabolism	41	45	0.0000963
	Biosynthesis of secondary metabolites	403	995	0.000518831
	Carotenoid biosynthesis	23	29	0.033348732
	Pentose phosphate pathway	34	54	0.039630093
A versus CK	Thiamine metabolism	5	11	0.008572893
	Glutathione metabolism	12	93	0.008572893
	Plant-pathogen interaction	16	164	0.010432522
	Sesquiterpenoid and triterpenoid biosynthesis	5	23	0.045374687
A versus Acy	Circadian rhythm - plant	26	36	0.00000000707
	Photosynthesis	29	77	0.0000237
	Porphyrin and chlorophyll metabolism	17	45	0.003560979
	Metabolic pathways	258	1,861	0.014721023
	Galactose metabolism	16	55	0.035306697
	Flavonoid biosynthesis	9	21	0.038121712
AcyA versus CK	Metabolic pathways	737	1,861	0.002207642
	Porphyrin and chlorophyll metabolism	38	45	0.002207642
	Carotenoid biosynthesis	26	29	0.012733735
	Biosynthesis of secondary metabolites	405	995	0.016208134
AcyA versus Acy	Flavonoid biosynthesis	14	21	0.0000000000895
	Brassinosteroid biosynthesis	5	8	0.001335017
	Zeatin biosynthesis	6	23	0.006857478
	Plant hormone signal transduction	19	271	0.048334673

Table 6. Continued.

Comparison Combination*	Term	Input No.	Background No.	Corrected P Value
AcyA versus A	Circadian rhythm - plant	25	36	0.0000000243
	Glyoxylate and dicarboxylate metabolism	20	63	0.008610509
	Nitrogen metabolism	15	42	0.013363167
	Porphyrin and chlorophyll metabolism	15	45	0.018115529

* A, *Aphis gossypii*; Acy, *Acyrtosiphon gossypii*; AcyA, *Acyrtosiphon gossypii* and *Aphis gossypii*; CK, XXX.

Hettenhausen et al. (2016) demonstrated that feeding by *Spodoptera exigua* Hübner or *Aphis glycines* Matsumura increased calcium-dependent protein kinase transcription in soybean, *Glycine max* (L.) Merrill. Sytykiewicz (2016) reported that feeding by *Rhopalosiphum padi* (L.) and *Sitobion avenae* (F.) significantly affected expression of *rbohA* and *rbohD* in maize, *Zea mays* L. In thale cress, *Arabidopsis thaliana* (L.) Heynh., feeding by *Bemisia tabaci* (Gennadius) biotype B nymphs induced the upstream jasmonic acid (JA) response genes *LOX2* and *OPR3* and inhibited the downstream JA response gene *VSP1*, while feeding by adults significantly inhibited the expression of *LOX2* and *OPR3* (Zhang et al. 2013a, b). In rice (*Oryza sativa* L.), feeding by either chewing or sucking insects affected the ethylene and JA pathways, and *OsHI-LOX* was a key gene in JA synthesis (Ma et al. 2020, Zhou et al. 2009). Aphids feeding on tobacco, *Nicotiana tabacum* L., foliage induced significantly fewer differentially expressed genes compared to feeding by mirids, mealybugs, or lepidopteran larvae (Heidel and Baldwin 2004). Our results showing differential expression of genes and the occurrence of enriched metabolic pathways in cotton after feeding by *Ac. gossypii* and *A. gossypii* further support those findings and demonstrate their involvement in the host plant defense response in cotton.

Physiological metabolic pathways are important regulatory pathways for plants to initiate defense responses. Related metabolic pathways participate in and complement various defense mechanisms such as local defense, systemic defense, and direct defense of plants (Dicke and Poেকে 2002, Orians 2005, Vignutelli et al. 1998). For example, *Apolygus lucorum* Meyer-Dur feeding induced significant changes in flavonoids, phenols, chymotrypsin inhibitors, condensed tannins, and amino acids in grape (*Vitis* spp.) leaves (Gao et al. 2019). Levels of soluble sugar, soluble protein, and chlorophyll in leaves of *Mikania micrantha* Kunth increased, while the activity of CAT, superoxide dismutase, and POD decreased after feeding by *Pachypeltis* sp. (Li et al. 2018). *Sitobion avenae* feeding on cabbage, *Brassica oleracea* (L.), and wheat, *Triticum aestivum* L., could induce increases in PPO, POD, and phenylalanine ammonia lyase activity (Han et al. 2009, Zhang et al. 2005). These studies show that piercing-sucking insect herbivory can cause changes in the reactive oxygen species system, secondary metabolite

synthesis, and other physiological metabolic pathways in host plants. Our results in this present study demonstrated that feeding by the aphids *Ac. gossypii* and *A. gossypii* also induced changes in multiple physiological metabolic pathways in cotton. These included photosynthetic and secondary metabolic pathways which could improve the cotton plant ability to compensate for damage or loss of photosynthates or other nutrients.

Our findings also showed that feeding by *Ac. gossypii* and *A. gossypii* on cotton significantly affected the functional expression of oxidoreductase enzymes in the host plant. The oxidoreductase system, including oxidoreductase lipooxygenase, propylene oxide synthase, propylene oxide cyclase, peroxidase, and polyphenol oxidase, is reported as an important protective enzyme system in defense reactions in cotton (Chung et al. 2013, Si et al. 2020, Ximénez-Embún et al. 2017). We further postulate that cotton initiates oxidoreductase gene expression immediately upon incurring pest damage, thus enhancing the host plant resistance to or tolerance of aphid feeding by increasing the level of protective enzymes (Yan et al. 2013, Zhang et al. 2020).

In addition, we found that when fed upon by either *Ac. gossypii* or *A. gossypii*, expression of photosynthesis-related genes was increased in cotton, which supports the findings of Gutsche et al. (2009) that insect feeding can upregulate the expression of photosynthesis-related genes in plants and the conclusion of Kangasjarvi et al. (2012) that photosynthesis is involved in plant defense responses as well as plant physiological functions as a remedy for carbon loss. Furthermore, feeding by combined populations of *Ac. gossypii* and *A. gossypii* on cotton significantly affected plant biological processes (e.g., cotton phosphorescence signal transduction). These results provide insight into mechanisms underlying the observed increase in chlorophyll and carotenoid content in cotton leaves when cotton is damaged by *Ac. gossypii* and *A. gossypii* (Deng et al. 2013, Zhang et al. 2020).

In our KEGG enrichment analysis, *Ac. gossypii* increased the expression of biosynthetic pathways of secondary metabolites in cotton, while feeding by *A. gossypii* increased the expression of sesquiterpenes and triterpenoids. When the two species fed together on the same plant, the expression of flavonoid biosynthesis genes increased. Therefore, under the stress of *Ac. gossypii* and *A. gossypii*, cotton initiates defense responses through different pathways involving secondary metabolism. It is known that plant metabolites, including flavonoids, terpenoids, alkaloids, and other secondary metabolites, play an important role in insect feeding induction which, when ingested by the insect, can inhibit digestion, affect feeding, or even kill the insect (Chen et al. 2019, Howe and Jander 2008). Previous studies have shown that *Ac. gossypii* and *A. gossypii* feeding increased levels of tannins, flavonoids, total phenols, and other secondary substances in cotton (Liu and Yang 1993, Wu 2020, Zhang 2020) and increased the activity of secondary metabolic enzymes in cotton (Li et al. 1998b, Lu et al. 2017).

When plants are fed upon by insects, they not only synthesize secondary metabolites that are toxic and deterrent, but they also produce changes in primary metabolites such as proteins and soluble sugars (Sulpice and McKeown 2015, Sun et al. 2013). Cotton plants fed upon by *Ac. gossypii* and *A. gossypii* respond by increasing soluble protein and sugar content as a defensive mechanism (Deng et al. 2013, Patima et al. 2018, Yan et al. 2013, Zhang et al. 2020), which corresponds

to an acceleration of biosynthesis and biological metabolism. Our GO and KEGG enrichment analyses indicated that steroid biosynthesis, lipid biosynthesis, and the pentose phosphate pathway were enhanced in cotton after feeding by *Ac. gossypii*. Those analyses also showed that cotton on fed by *A. gossypii* exhibited enhanced methionine metabolism, and that feeding by mixed populations of the aphids enhanced the differential expression of galactose metabolism.

These cotton plant reactions to aphid attack are defense mechanisms. Sterols involved in steroid biosynthesis, lipid biosynthesis, and methionine metabolism play an important role in cell wall formation, cell elongation, and development (Carland et al. 2002, Catterou et al. 2001, Clouse and Sasse 1998, He et al. 2003), while methionine is directly involved in protein biosynthesis (Giovannelli et al. 1985). Metabolites from galactose metabolism can promote the cell wall formation (Atmodjo et al. 2013) and increase the content of soluble sugars in plants (Thoden and Holden 2005). These plant defense mechanisms are energy-consuming processes (Coley et al. 1985, Mooney and Gulmon 1982, Rhoades 1979), and ATP is continuously provided for these processes through the pentose phosphate pathway. Collectively, these biosynthetic processes strongly influence plant morphology, protein and carbohydrate synthesis, and continuous plant defense functions (Limdsey et al. 2003, Schaller 2003).

Molecular studies of plant-pest interactions can reveal crop insect resistance mechanisms. Antibiotic-related substances, such as disease-related proteins, are rapidly produced when plants are fed upon by sucking insects (Park et al. 2005). Furthermore, oxygen-burst reactions occur at injured sites of plants, resulting in accumulation of protein I proteins and injury responses (Kaloshian 2004), activation of mitogen-activated protein kinases, synthesis and interaction of phytohormones, and a series of stress responses in plants (Erb et al. 2012, Zebelo and Maffei 2015). Aphids may also transmit viral plant diseases while feeding (Feres and Moreno 2009). Cotton will immediately initiate immune factors to resist viral infection (Körner et al. 2013, Mandadi and Scholthof 2013). In support, our KEGG enrichment analysis showed that the largest number of differentially expressed genes were annotated to the plant-pathogen interaction pathway in cotton fed upon by the aphids.

It should be noted that when *A. gossypii* and *Ac. gossypii* feed on cotton in mixed populations, the genes regulating circadian rhythm are differentially expressed, which may be related to the regulation of nutrient homeostasis (Haydon et al. 2015), hormone synthesis and signal transduction (Atamian and Harmer 2016), redox reaction (Zhou et al. 2015), and the changes in levels of some major osmotic regulators (Greenham and McClung 2015). These responses indicate that the biological clock of cotton has a complex regulation when stressed by aphid feeding.

In conclusion, although the gene expression and metabolic pathways of cotton defense responses induced by *A. gossypii* and *Ac. gossypii* differ, they all enhance the defense response of cotton through regulating pathways related to photosynthetic substances, oxidoreductase activity, secondary metabolism, and other metabolic activities. This is similar to the defense response pathways induced by most insects with piercing-sucking mouthparts and feeding habits. When the two aphid species damaged the plant simultaneously, the genes regulating cotton photosynthetic phosphorus signal transduction, circadian rhythm regulation, porphyrin and chlorophyll metabolism, photosynthesis, galactose metabolism,

flavonoid biosynthesis, and other activities were significantly expressed. Our study thus provides new insights into the complex mechanisms underlying cotton defense responses against aphid attacks. However, in this study, only single omics analysis was used to analyze the mechanisms of cotton defense against aphids. In the future, multigroup analysis should be used to conduct more in-depth analysis at the molecular, metabolic, and protein levels, so as to provide a more comprehensive elucidation of the mechanism of cotton defense against aphids.

Acknowledgment

This work was supported by the National Natural Science Foundation of China (grant no. 31660519).

References Cited

- Anders, S. 2010.** HTSeq: Analysing high-throughput sequencing data with Python. European Molecular Biology Laboratory. https://htseq.readthedocs.io/en/release_0.9.1/ (Last accessed 12 October 2019).
- Anders, S. and W. Huber. 2012.** Differential expression of RNA-Seq data at the gene level—The DESeq package. European Molecular Biology Laboratory. https://htseq.readthedocs.io/en/release_0.9.1/ (Last accessed 12 October 2019).
- Atamian, H.S. and S.L. Harmer. 2016.** Circadian regulation of hormone signaling and plant physiology. *Plant Mol. Biol.* 91: 691–702.
- Atmodjo, M.A., Z. Hao and D. Mohnen. 2013.** Evolving views of pectin biosynthesis. *Annu. Rev. Plant Biol.* 64: 747–779.
- Boyko, E.V., C.M. Smith, V.K. Thara, J.M. Bruno, Y. Deng, S.R. Starkey and D.L. Klahsen. 2006.** Molecular basis of plant gene expression during aphid invasion: Wheat Pto- and Pti-like sequences are involved in interactions between wheat and Russian wheat aphid (Homoptera: Aphididae). *J. Econ. Entomol.* 99: 1430–1445.
- Bricchi, I., C.M. Berteà, A. Occhipinti, L.A. Paponov and M.E. Maffei. 2012.** Dynamics of membrane potential variation and gene expression induced by *Spodoptera littoralis*, *Myzus persicae*, and *Pseudomonas syringae* in *Arabidopsis*. *PLoS One* 7: e46673.
- Carland, F.M., S. Fujioka, S. Takatsuto, S. Yoshida and T. Nelson. 2002.** The identification of CVP1 reveals a role for sterols in vascular patterning. *Plant Cell* 14: 2045–2058.
- Catterou, M., F. Dubois, H. Schaller, L. Aubanelle, B. Vilcot, B.S. Sangwan-Norreel and R.S. Sangwan. 2001.** Brassinosteroids, microtubules and cell elongation in *Arabidopsis thaliana*. I. Molecular, cellular and physiological characterization of the *Arabidopsis bul* mutant, defective in the Δ^7 -sterol-C5-desaturation step leading to brassinosteroid biosynthesis. *Planta* 212: 659–672.
- Chen, L.-H., W. Patima, Y.-H. Cui, Y. Li and H.Z. Feng. 2015.** Effect of *Aphis gossypii* feeding on defense enzyme activity in different varieties of cotton. *Xinjiang Agric. Sci.* 52: 1866–1871.
- Chen, X., D.-D. Wang, X. Fang, X.-Y. Chen and Y.-B. Mao. 2019.** Plant specialized metabolism regulated by jasmonate signaling. *Plant Cell Physiol.* 60: 2638–2647.
- Chen, Y.-C., C.-X. Cao, Z.-J. Huang, T. Wang, C.-M. Ji and G.-H. Lei. 2011.** Study on the nutrient materials and the resistance of cucumber aphid. *Chin. Agric. Sci. Bull.* 27: 283–386.
- Chung, S.H., C. Rosa, E.D. Scully, M. Peiffera, J.F. Tookera, K. Hoovera, D.S. Luthe and G.W. Feltona. 2013.** Herbivore exploits orally secreted bacteria to suppress plant defenses. *Proc. Natl. Acad. Sci. U. S. A.* 110: 15728–15733.
- Clouse, S.D. and J.M. Sasse. 1998.** Brassinosteroids: Essential regulators of plant growth and development. *Plant Physiol.* 49: 472–451.

- Coley, P.D., J.P. Bryant and F.S. Chapin. 1985. Resource availability and plant antiherbivore defense. *Science* 230: 895–899.
- Deng X.-X., H.-L. Jiang, J. Peng, Z.-M. He, T.-W. Ma and J.-G. Wang. 2013. Physiological responses of cotton to feeding by *Aphis gossypii* during the flower-bolling stage. *Chin. J. Appl. Entomol.* 50: 161–166.
- Dicke, M. and R.M.P. v. Poecke. 2002. Signaling in plant-insect interactions: Signal transduction in direct and indirect plant defence, Pp. 289–316. In Scheel, D. and C. Wasternack (eds.), *Plant Signal Transduction*. Oxford Univ. Press, Oxford, England.
- Erb, M., S. Meldau and G.A. Howe. 2012. Role of phytohormones in insect specific plant reactions. *Trends Plant Sci.* 17: 250–159.
- Fereres, A. and A. Moreno. 2009. Behavioural aspects influencing plant virus transmission by homopteran insects. *Virus Res.* 141: 158–168.
- Gao, J.-M., S.-H. Gao and B.-J. Gao. 2019. Effects of insect feeding and mechanical damages on metabolites in leaves of grape. *Acta Agric. Bor.-Occid. Sin.* 28: 1543–1551.
- Giovanelli, J., M.S. Harvey and H.D. Anne. 1985. Quantitative analysis of pathways of methionine metabolism and their regulation in Lemna. *Plant Physiol.* 78: 555–560.
- Greenham, K. and C.R. McClung. 2015. Integrating circadian dynamics with physiological processes in plants. *Nat. Rev. Genet.* 16: 598–610.
- Gutsche, A., T. Heng-Moss, G. Sarath, P. Twigg, Y. Xia, G. Lu and D. Mornhinweg. 2009. Gene expression profiling of tolerant barley in response to *Diuraphis noxia* (Hemiptera: Aphididae) feeding. *B. Entomol. Res.* 99: 163–173.
- Han, Y., Y. Wang, J.-L. Bi, X.-Q. Yang, Y. Huang, X. Zhao, Y. Hu and Q.-N. Cai. 2009. Constitutive and induced activities of defense-related enzymes in aphid-resistant and aphid-susceptible cultivars of wheat. *J. Chem. Ecol.* 35: 176–182.
- Haydon, M.J., Á. Román and W. Arshad. 2015. Nutrient homeostasis within the plant circadian network. *Front Plant Sci.* 6: 299.
- He, J.-X., S. Fujioka, T.-C. Li, S.-G. Kang, H. Seto, S. Takatsuto, S. Yoshida and J.-C. Jang. 2003. Sterols regulate development and gene expression in Arabidopsis. *Plant Physiol.* 131: 1258–1269.
- Heidel, A.J. and I.T. Baldwin. 2004. Microarray analysis of salicylic acid- and jasmonic acid-signalling in responses of *Nicotiana attenuata* to attack by insects from multiple feeding guilds. *Plant Cell Environ.* 27: 1362–1373.
- Hettenhausen, C., G. Sun, Y. He, H. Zhuang, T. Sun, J. Qi and J. Wu. 2016. Genome-wide identification of calcium-dependent protein kinases in soybean and analyses of their transcriptional responses to insect herbivory and drought stress. *Sci. Rep.* 6: 18973.
- Howe, G.A. and G. Jander. 2008. Plant immunity to insect herbivores. *Annu. Rev. Plant Biol.* 59: 41–66.
- Hullé, M., B. Chaubet, E. Turpeau and J.-C. Simon. 2020. Encyclop'Aphid: A website on aphids and their natural enemies. *Entomol. Gen.* 40: 97–101.
- Jacobson, R. and P. Croft. 1998. Strategies for the control of *Aphis gossypii* Glover (Hom.: Aphididae) with *Aphidius colemani* Viereck (Hym.: Braconidae) in protected cucumbers. *Biocontr. Sci. Tech.* 8: 377–387.
- Kaloshian, I. 2004. Gene-for-gene disease resistance: bridging insect pest and pathogen defence. *J. Chem. Ecol.* 30: 2419–2438.
- Kanehisa, M., M. Araki, S. Goto, M. Hattori, M. Hirakawa, M. Itoh, T. Katayama, S. Kawashima, S. Okuda, T. Tokimatsu and Y. Yamanishi. 2008. KEGG for linking genomes to life and the environment. *Nucleic Acids Res.* 36: 480–484.
- Kangasjarvi, S., J. Neukermans, S. Li, E.-M. Aro and G. Noctor. 2012. Photosynthesis, photorespiration, and light signalling in defence responses. *J. Exp. Bot.* 63: 1619–1636.
- Kim, D., B. Langmead and S.L. Salzberg. 2015. HISAT: A fast spliced aligner with low memory requirements. *Nat. Methods* 12: 357–360.
- Körner, C. J., D. Klauser, A. Niehl, A. Domínguez-Ferreras, D. Chinchilla, T. Boller, M. Heinlein and D.R. Hann. 2013. The immunity regulator BAK1 contributes to resistance against diverse RNA viruses. *Mol. Plant Microbe Interact.* 26: 1271–1280.

- Li, C.-X., L.-F. Gao, X. Mao and R.-Z. Li. 1998a.** Responses of cotton antioxidant system to aphid feeding suppression. *J. Shanxi Agric. Univ.* 18: 207–210.
- Li, R.-Z., X. Mao, C.-X. Li and L.-F. Gao. 1998b.** The relationships between activities of secondary metabolism-related enzymes and induced resistance to aphids in cotton. *J. Shanxi Agric. Univ.* 18: 165–168.
- Li, S., D.-J. Zheng, Y. Wang, Y. Lin, M.-R. Liu, F.-R. Gui, F.-H. Wan and C. Xi. 2018.** Effects of feeding by *Pachypeltis* sp. (Hemiptera: Miridae) on the defensive enzymes and nutrients in *Mikania micrantha* leaves. *J. Biosafety* 27: 45–49.
- Limdsey, K., M.L. Pullen and J.F. Topping. 2003.** Importance of plant sterols in pattern formation and hormone signalling. *Trends Plant Sci.* 8: 521–525.
- Liu, C.-Z. and J.-N. Lan. 2009.** Variations of oxidase in the seedling of three alfalfa varieties infested by *Therioaphis trifolii* Monell (Homoptera: Aphididae). *Acta Agrestia Sin.* 17: 32–35.
- Liu, X.-M. and Q.-H. Yang. 1993.** The relationships between the physiological and biochemical mechanisms of aphid resistance of cotton and the population dynamics of cotton aphid. *Acta Phyto. Sin.* 20: 27–29.
- Lu, Y.-H. and G.-M. Liang. 2016.** Research advance on the succession of insect pest complex in Bt crop ecosystem. *Plant Protect.* 24: 7–11.
- Lu, Y.-H., G.-M. Liang, Y.-J. Zhang and X.-M. Yang. 2020.** Advances in the management of insect pests of cotton in China since the 21st century. *Chin. J. App. Entomol.* 57: 477–490.
- Lu, Y., H.-Q. Wang, L.-S. Chen, P.-L. Wang and J. Li. 2017.** Correlation between induced resistance to aphids and secondary metabolism enzyme activities of cotton varieties in Xinjiang. *Plant Protect.* 43: 51–55.
- Luo, S.-T., X. Zhang, J.-F. Wang, C.-Y. Jiao, Y.-Y. Chen and Y.-B. Shen. 2017.** Plant ion channels and transporters in herbivory-induced signalling. *Funct. Plant Biol.* 45: 111–131.
- Luo, Z., H.-Z. Dong, X.-Q. Kong, Z. Ming and Y.-Q. Zhu. 2008.** Individual and combined effects of salinity and waterlogging on Cry1Ac expression and insecticidal efficacy of Bt cotton. *Crop Prot.* 27: 1485–1490.
- Ma, F.-L., X.-F. Yang, Z.-Y. Shi and X.-X. Miao. 2020.** Novel crosstalk between ethylene- and jasmonic acid-pathway responses to a piercing-sucking insect in rice. *New Phytol.* 225: 474–487.
- Mandadi, K.K. and K.-B.G. Scholthof. 2013.** Plant immune responses against viruses: How does a virus cause disease? *Plant Cell* 25: 1489–1505.
- Martinez-Medina, A., V. Flors, M. Heil, B. Mauch-Mani, C.M.J. Pieterse, M.J. Pozo, J. Ton, N.M. van Dam and U. Conrath. 2016.** Recognizing plant defense priming. *Trends Plant Sci.* 21: 818–822.
- Mooney, H.A. and S.L. Gulmon. 1982.** Constraints on leaf structure and function in reference to herbivory. *Bioscience* 32: 198–206.
- Moran, P.J. and G.A. Thompson. 2001.** Molecular responses to aphid feeding in arabidopsis in relation to plant defense pathways. *Plant Physiol.* 125: 1074–1085.
- Mortazavi, A., B. A. Williams, K. McCue, L. Schaeffer and B. Wold. 2008.** Mapping and quantifying mammalian transcriptomes by RNA-Seq. *Nat. Methods* 5: 621–628.
- Orians, C. 2005.** Herbivores, vascular pathways, and systemic induction: Facts and artifacts. *J. Chem. Ecol.* 31: 2231–2242.
- Park, S.J., Y. Huang and P. Ayoubi. 2005.** Identification of expression profiles of sorghum genes in response to greenbug phloem-feeding using cDNA subtraction and microarray analysis. *Planta* 223: 932–947.
- Patima W., S.-J. Ma, P.-P. Guo, M.-M. Wu, F. Liu and D.-Y. Ma. 2018.** Effects of *Aphis gossypii* feeding stresses with different densities on soluble sugar and free proline contents in cotton leaves. *J. Xinjiang Agric. Univ.* 41: 117–120.
- Rhoades, D.F. 1979.** Evolution of plant chemical defense against herbivores, Pp. 4–54. *In* Rosenthal, G.A. and Janzen, D. H. (eds.), *Herbivores: Their Interaction With Secondary Plant Metabolites*. Academic Press, New York.

- Robinson, M.D., D.J. McCarthy and G.K. Smyth. 2010.** edgeR: A bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* 26: 139–140.
- Schaller, H. 2003.** The role of sterols in plant growth and development. *Prog. Lipid Res.* 42: 163–175.
- Si, H., H.-L. Liu, Y.-W. Sun, Z.-P. Xu, S.-J. Liang, B. Li, X. Ding, J.-Y. Li, Q.-Q. Wang, L. Sun, X.-L. Zhang and S.-X. Jin. 2020.** Transcriptome and metabolome analysis reveal that oral secretions from *Helicoverpa armigera* and *Spodoptera litura* influence wound-induced host response in cotton. *Crop J.* 8: 929–942.
- Sulpice, R. and P.C. McKeown. 2015.** Moving toward a comprehensive map of central plant metabolism. *Annu. Rev. Plant Biol.* 66: 187–210.
- Sun, J.-R., H.-J. Yang, T. Zhang, C.-J. Cao, S.-X. Zong, Y.-Q. Luo and Y.-B. Shen. 2013.** Metabolites of *Ammopiptanthus mongolicus* induced by *Orgyia ericae* attack and mechanical wounding. *Plant Physiol. Bioch.* 69: 101–107.
- Sytykiewicz, H. 2016.** Deciphering the role of NADPH oxidase in complex interactions between maize (*Zea mays* L.) genotypes and cereal aphids. *Biochem. Bioph. Res. Co.* 476: 90–95.
- Thoden, J.B. and H.M. Holden. 2005.** The molecular architecture of galactose: Mutarotase/UDP-galactose 4-epimerase from *Saccharomyces cerevisiae*. *J. Biol. Chem.* 280: 21900–21907.
- Thompson, G.A. and F.L. Goggin. 2006.** Transcriptomics and functional genomics of plant defence induction by phloem-feeding insects. *J. Exp. Bot.* 57: 755–766.
- Tjallingii, W.-F. 2006.** Salivary secretions by aphids interacting with proteins of phloem wound responses. *J. Exp. Bot.* 57: 739–745.
- Vignutelli, A., C. Wasternack, K. Apel and H. Bohlmann. 1998.** Systemic and local induction of an *Arabidopsis thionin* gene by wounding and pathogens. *Plant J.* 14: 285–295.
- Vincent, T.R., M. Avramova, J. Canham, J. Canham, P. Higgins, N. Bilkey, S.T. Mugford, M. Pitino, M. Toyota, S. Gilroy, A.J. Miller, S. Hogenhout and D. Sanders. 2017.** Interplay of plasma membrane and vacuolar ion channels, together with BAK1, elicits rapid cytosolic calcium elevations in *Arabidopsis* during aphid feeding. *Plant Cell* 29: 1460–1479.
- Voelckel, C., W.W. Weisser and I.T. Baldwin. 2004.** An analysis of plant–aphid interactions by different microarray hybridization strategies. *Mol. Ecol.* 13: 3187–3195.
- Wang, H.-T., M. Xue, Q.-L. Li, X. Zhang and F.-Y. Zhou. 2011.** Effects of tobacco plants infested by *Bemisia tabaci* (Gennadius) B-biotype on the development and reproduction of *Spodoptera litura* (Fabricius) and related mechanisms. *Sci. Agric. Sin.* 44: 4600–4609.
- Wang, L.-K., Z.-X. Feng, X. Wang, X.-W. Wang and X.-G. Zhang. 2010.** DEGseq: An R package for identifying differentially expressed genes from RNA-seq data. *Bioinformatics* 26: 136–138.
- Wu, M.-M, L.-R. Yang, X.-Z. Yang, X. Zhang, J.-X. Ma, Y. Zhang and H. Li. 2015.** Effect of aphid invasion on physiological and biochemical indexes in watermelon seedling stage. *J. Fruit Sci.* 32: 943–949.
- Wu, N. 2020.** Studies on the physiological metabolism of *Aphis gossypii* and *Acyrtosiphon gossypii* induced by interactive feeding on cotton. M.S. Thesis, Shihezi Univ. Shihezi, China.
- Ximénez-Embún, M.G., J.J. Glas, F. Ortego, J.M. Alba, P. Castañera and M.R. Kant. 2017.** Drought stress promotes the colonization success of a herbivorous mite that manipulates plant defenses. *Exp. Appl. Acarol.* 73: 297–315.
- Yan, C., M. Fan, M. Yang, J.-P. Zhao, W.-H. Zhang, Y. Su, L.-T. Xiao, H.-T. Deng and D.-X. Xie. 2018.** Injury activates Ca^{2+} /calmodulin-dependent phosphorylation of JAV1-JAZ8-WRKY51 complex for jasmonate biosynthesis. *Mol. Cell* 70: 136–149.
- Yan, H.-M., X.-X. Deng, F.-J. Wang, Y. Gao, Q.-L. Wang, H.-L. Jiang and J.-G. Wang. 2013.** Effect of different aphid damage levels on the physiology and biochemistry of bud stage cotton. *Xinjiang Agric. Sci.* 50: 2077–2084.

- Young, M.D., M.J. Wakefield, G.K. Smyth and A. Oshlack. 2010.** Gene ontology analysis for RNA-seq: Accounting for selection bias. *Genome Biol.* 11: R14.
- Zebelo, S.A. and M.E. Maffei. 2015.** Role of early signalling events in plant insect interactions. *J. Exp. Bot.* 66: 435–448.
- Zhang, C.-N., J.-X. Wu, W. Dai and L. Chen. 2005.** Activities of some isoenzymes in the leaves of *Brassica oleracea* seedlings infested by peach aphid (*Myzus persicae*). *Acta Bot. Bor.-Occid. Sin.* 25: 1566–1569.
- Zhang, P.-J., W.-D. Li, F. Huang, J.-M. Zhang, F.-C. Xu and Y.-B. Lu. 2013a.** Feeding by whiteflies suppresses downstream jasmonic acid signaling by eliciting salicylic acid signaling. *J. Chem. Ecol.* 39: 612–619.
- Zhang, P.-J., C.-X. Xu, J.-M. Zhang, Y.-B. Lu, J.-N. Wei, Y.-Q. Liu, A. David, W. Boland and T.C.J. Turlings. 2013b.** Phloem-feeding whiteflies can fool their host plants, but not their parasitoids. *Funct. Ecol.* 27: 1304–1312.
- Zhang, S.-F., X.-B. Kong, H.-B. Wang and Z. Zhang. 2013c.** Different strategies of plant resistance to insects and their interactions. *Chin. J. App. Entomol.* 50: 1428–1437.
- Zhang, Y.-D. 2020.** The cotton defense response induced by *Acyrtosiphon gossypii* and its effect on late feeding of *Aphis gossypii*. M.S. Thesis, Shihezi University. Shihezi, China.
- Zhang, Y.-D., N. Wu, X.-H. Cai, Y.-H. Shi and J.-G. Wang. 2020.** Effects of *Acyrtosiphon gossypii* on physiology, biochemistry and related defense enzymes of cotton. *Xinjiang Agric. Sci.* 57: 2065–2074.
- Zhou, G.-X., J.-F. Qi, N. Ren, J.-A. Cheng, M. Erb, B.-Z. Mao and Y.-G. Lou. 2009.** Silencing Os HI-LOX makes rice more susceptible to chewing herbivores, but enhances resistance to a phloem-feeder. *Plant J.* 60: 638–648.
- Zhou, M., W. Wang, S. Karapetyan, M. Mwimba, J. Marqués, N.E. Buchler and X.-N. Dong. 2015.** Redox rhythm reinforces the circadian clock to gate immune response. *Nature* 523: 472–476.