# Effectiveness and Biochemical Impact of Flubendiamide and Flonicamid Insecticides against *Bemisia tabaci* (Hemiptera: Aleyrodidae) and Residue Dissipation in Cherry Tomato Plants and Soil under Greenhouse Conditions<sup>1</sup>

Moataz A.M. Moustafa<sup>2</sup>, Dalia E. El Hefny<sup>3</sup>, Nawal Abdulaziz Alfuhaid<sup>4</sup>, Rania M.A. Helmy<sup>3</sup>, Nourhan A. El-Said, and El-Desoky S. Ibrahim

Department of Economic Entomology and Pesticides, Faculty of Agriculture, Cairo University, 12613 Giza, Egypt

Abstract We investigated the effectiveness and biochemical impact of the insecticides flubendiamide and flonicamid compared with azadirachtin and acetamiprid against the whitefly Bemisia tabaci (Grennadius) (Hemiptera: Aleyrodidae) infesting cherry tomato (Solanum lycopersicum var. cerasiforme L.) grown under greenhouse conditions. The dissipation of both insecticides in the plants and in the soil also was determined using the QuEChERS (quick, easy, cheap, effective, rugged, and safe) and liquid chromatography-electrospray ionization-mass spectrometry methods. Both insecticides were more effective for reducing B. tabaci populations than were either acetamiprid or azadirachtin. Biochemical analysis revealed that esterase may play an important role in whitefly adaptation to flubendiamide and flonicamid. The QuEChERS method was determined suitable for quickly detecting residues of flubendiamide and flonicamid in complex matrices. The recovery rates on tomato fruit samples were 92.8-106.0%, with a relative standard deviation (RSD) range of 0.46-2.65%. For soil samples, the recovery rates were 81.3-95.7% with RSDs of 1.20-3.86%. We further determined that flubendiamide had dissipation half-lives of 3.13, 3.63, and 3.68 d in tomato fruit, tomato leaves, and soil, respectively. Flonicamid had half-lives of 4.25, 3.54, and 2.60 d in fruit, leaves, and soil, respectively. These results suggest that preharvest intervals of 3 and 7 d are appropriate for flubendiamide and flonicamid, respectively, in cherry tomato production. The risk quotient was >1 by the day 5 after application; however, that value declined to <1 on day 7 after application, indicating little long-term risk to human health.

Key Words flubendiamide, flonicamid, Bemisia tabaci, enzymes activity, dissipation

Cherry tomato (Solanum lycopersicum var. cerasiforme L.) production in greenhouses is compromised by the whitefly *Bemisia tabaci* Gennadius (Hemiptera: Aleyrodidae), causing significant yield losses sometimes as high as 100%

J. Entomol. Sci. 59(3): 289–310 (July 2024) DOI: 10.18474/JES23-61

<sup>&</sup>lt;sup>1</sup>Received 6 September 2023; accepted for publication 14 October 2023.

<sup>&</sup>lt;sup>2</sup>Corresponding author (email: moataz.moustafa79@gmail.com).

<sup>&</sup>lt;sup>3</sup>Pesticide Residues and Environmental Pollution Department, Central Agricultural Pesticide Laboratory, Agricultural Research Center, Dokki, 12618 Giza, Egypt.

<sup>&</sup>lt;sup>4</sup>Department of Biology, College of Science and Humanities, Prince Sattam Bin Abdulziz University, Al-Kharj 11942, Saudi Arabia.

(McKenzie et al. 2014, Mugerwa et al. 2021, Ochilo et al. 2019). Moodley et al. (2019) estimated the losses as exceeding US\$100 million each year. Therefore, conventional chemical insecticides are routinely used as a preferred tactic for preventing and managing whitefly attacks on tomatoes, thus ensuring tomato quality and yield (Ghosal and Chatterjee 2018). The performance of novel ecofriendly insecticides against *B. tabaci* and other whitefly pests is continually being evaluated (Kumar et al. 2019).

Synthetic amide insecticides are a class of novel insecticides that have excellent effectiveness against target insects and low toxicity for nontarget species (Lin et al. 2021). Flubendiamide is a diamide insecticide that was developed by Bayer Crop Science (Leverkusen, Germany) and has been used recently (Casida 2015, Sparks and Nauen 2015) against many lepidopteran, dipteran, and coleopteran insects (Kadala et al. 2020, Li et al. 2019). It has a novel mode of action, acting as an activator of the insect ryanodine receptor and causing massive release of calcium ions from muscle cells (Cordova et al. 2006, Uesugi et al. 2020). However, because of its environmental behavior and ecological toxicity, flubendiamide may be a danger to invertebrates and poses a significant risk to aquatic habitats (Lin et al. 2021).

Flonicamid (*N*-(cyanomethyl)-4-(trifluoromethyl)-3-pyridinecarbox-amide) is a new selective systemic insecticide with extremely potent insecticidal efficacy against whiteflies and other piercing-sucking insects (Morita et al. 2007). It works by obstructing type-A potassium channels, thus preventing insects from moving to and attacking tomato plants (Morita et al. 2000).

Because pesticide residues may adversely affect the quality and growth of tomato plants, the QuEChERS (quick, easy, cheap, effective, rugged, and safe) method is currently used for the rapid identification of pesticides in agricultural goods (Anastassiades et al. 2003). Standard QuEChERS methods also have been used for clean-up procedures with primary secondary amine (PSA), clean-up C18 bulk sorbent, and graphitized carbon black (GCB) as adsorbents (Anastassiades et al. 2003, El-Hefny et al. 2021). Those chemicals effectively absorb pigments and minimize matrix effects. The residual behavior, dissipation behavior, and dietary risk of tested pesticides in various crops also must be assessed to ensure safety and protect the environment (Zhang et al. 2022) by estimating the risk quotient, taking into consideration dissipation as an indicator for human health safety (El Hefny et al. 2021, Moustafa et al. 2023a).

In this study, the biological effectiveness and biochemical impact of flubendiamide and flonicamid insecticides, as alternatives to azadirachtin and acetamiprid, for managing whiteflies were assessed in developing a pest management spray schedule for cherry tomato crops. The dissipation of both insecticides also was determined in cherry tomato fruits and leaves and in soil using the QuEChERS and liquid chromatography–electrospray ionization tandem mass spectrometry (LC-ESI MS/MS) methods.

### Materials and Methods

**Insecticides and chemicals.** Four commercial insecticide formulations (azadirachtin; Gaara Co., Cairo, Egypt; acetamiprid, Shoura Chemicals Co., Giza, Egypt; flubendiamide, Sama-Trade Co., Giza, Egypt; and flonicamid, Shoura Chemicals Co., Giza, Egypt) were used in this study (Table 1). The flubendiamide

Common Name	Trade Name (% active ingredient)	Chemical Group	Application Rate (/100 L of water)
Azadirachtin	Ashouk (0.15% EC)	Biopesticide	350 ml
Acetamiprid	Mospilan (20% SP)	Neonicotinoid	25 g
Flubendiamide	Takumi (20% WG)	Diamide	50 g
Flonicamid	Tebekki (50% WG)	Pyridine carboxamide	50 g

Table 1. Tested insecticides and their application rates.

and flonicamid reference materials (98%) were purchased from Dr. Ehrenstorfer GmbH (Augsburg, Germany). Extraction and clean-up chemicals were obtained from Merck Company (Darmstadt, Germany). Anhydrous magnesium sulfate (MgSO<sub>4</sub>) fine powder and sodium chloride (NaCl) that had been dried in an oven at 250°C for 4 h were stored in desiccators prior to use. The PSA sorbent (Bondesil-PSA, 40 M) was obtained from Supelco Analytical Products (Sigma-Aldrich Company, Darmstadt, Germany). Stock solutions (1000 mg/L) of flubendiamide and flonicamid were prepared with acetonitrile. To prepare the sample matrices for high-performance liquid chromatography (HPLC) detection, calibrated matrix and solvent solutions at concentrations of 10, 5, 1, 0.5, 0.1, 0.05, and 0.01 mg/L were applied to the sample matrices. The resulting mixes were then passed through a 0.22-µm-pore size filter.

Greenhouse experiment. To evaluate the effectiveness of the tested insecticides against *B. tabaci* nymphs, greenhouse experiments were conducted at the Faculty of Agriculture (Giza, Egypt) over two consecutive seasons (2022 and 2023). Cherry tomato (cv. 'Katalina 522') seedlings with four or five true leaves were transplanted in a randomized complete block design with four replicates. The greenhouse (10  $\times$  60 m) was divided into five beds. Each bed was 1.2 m wide. The in-row distance between plants was 50 cm. Each experimental unit consisted of 10 plants. This area was managed with recommended agricultural practices for the entire season. The insecticides were applied with a Cooper Pegler  $CP_3$  knapsack sprayer with irrigation water used for dilutions (Moustafa et al. 2022). The chemical application began when the whitefly infestation reached four nymphs per leaf. After insecticide application, randomly selected tomato leaves were collected at 1, 3, 5, 7, 10, 15, and 21 d after application. One hundred leaves (25 leaves/replicate) were collected from each treatment for analysis (Moustafa et al. 2022, Kandil et al. 2023). The percentage of infestation reduction was calculated based on B. tabaci nymph infestation before and after application of each insecticide according to Henderson and Tilton (1955).

**Enzyme sample preparation.** Adults of *B. tabaci* were collected at 1, 7, 15, and 21 d after flubendiamide and flonicamid application at their recommended concentrations. Ten milligrams of adults from each treatment and control group were weighed and homogenized in phosphate buffer (0.1 M, pH 7). Three replicates were made for each treatment. The homogenates were centrifuged at 12,000  $\times$  *g* for 15 min, and the supernatant was stored at  $-20^{\circ}$ C.

Total esterase activity. Total esterase activity was analyzed with  $\alpha$ -naphthyl acetate ( $\alpha$ -NA) as a substrate, according to Van Asperen (1962) with some modifications (Moustafa et al. 2023b, 2023c). Thirty microliters of supernatant was mixed with 30 mM  $\alpha$ -NA for 15 min, after which Fast Blue B solution was added to stop the reaction. The absorbance was read at 600 nm, and the activity was assessed by comparison with the  $\alpha$ -naphthol standard curve.

**Glutathione S-transferase (GST) activity.** GST activity was measured according to Habing et al. (1974) and Moustafa et al. (2023b, 2023c). Ten microliters of enzyme solution was added to 25  $\mu$ l of a solution of 1-chloro-2,4-dinitrobenzene (30 mM) and 25  $\mu$ l of glutathione (50 mM dissolved in pH 6.5). The enzyme kinetics readings were recorded at 340 nm for 5 min.

**Cytochrome P-450 activity.** Cytochrome P-450 (a mixed function oxidase) activity was determined as described by Hansen and Hodgson (1971) and Moustafa et al. (2023b, 2023c) using *p*-nitrophenol to generate the standard curve. The substrate of *p*-nitroanisole (2 mM) was mixed with 90  $\mu$ l of enzyme solution for 2 min. Nicotinamide adenine dinucleotide phosphate (9.6 mM) was then added to the mixture. The absorbance was read at 405 nm for 15 min.

Statistical analysis of efficacy and enzyme activity. The effectiveness and biochemical activity of the tested insecticides were coded and entered using the statistical package SPSS (Statistical Package for the Social Sciences, v. 22, IBM, Armonk, NY). The results were tested for satisfying assumptions of parametric tests. The continuous variables were subjected to Shapiro-Wilk and Kolmogorov-Smirnov test for normality. The reduction percentage data were standardized for normality using the arcsine square root. Data are presented as mean  $\pm$  standard deviation. Analyses of variance were conducted for effectiveness and enzyme activity using MiniTab software (v. 14.0). The post hoc analysis was performed with a Tukey pairwise honestly significant difference (HSD) test, and results were considered significant at P < 0.05. The data were visualized, when possible, using R studio (v. 2022.02.4.).

**Residue analysis sample processing.** Purification and extraction were performed according to the methods of Anastassiadis et al. (2003) and Moustafa et al. (2023a). Cherry tomato fruit and leaf samples were collected randomly 2 h after spraying (0 d) and 1, 3, 5, 7, 10, 15, and 21 d after treatment. A homogenized tomato sample (10 g) was weighed into a 50-ml Teflon centrifuge tube to which 10 ml of acetonitrile was added. Each tube and its contents were shaken for 2 min on a Vortex mixer to ensure that the solvent interacted with the entire sample. One gram of NaCl and 4 g of MgSO<sub>4</sub> were then added to the solution and shaken for 1 min. A 10-ml Teflon centrifuge tube containing 50 mg of PSA and 300 mg of MgSO<sub>4</sub> for fruits and 50 mg of GCB for leaves was filled with 2 ml of the cleared supernatant following centrifugation at 5,000 rpm for 5 min. The mixture was then shaken for 1 min and centrifuged in a microcentrifuge for 5 min at 5,000 rpm.

Soil (5 g) was placed in a 50-ml Teflon centrifuge tube. To ensure that the solvent interacted with the entire sample, 10 ml of acetonitrile and 5 ml of water were added. The tube was then shaken for 2 min on a Vortex mixer. Following the addition of 1 g of NaCl and 4 g of MgSO<sub>4</sub>, the tube and its contents were again shaken for 1 min. A 10-ml Teflon centrifuge tube with 50 mg of PSA and 300 mg of MgSO<sub>4</sub> was filled with 2 ml of the cleared supernatant following centrifugation at 5,000

rpm for 5 min. After shaking for 1 min, the mixture was centrifuged with a microcentrifuge for 5 min at 5,000 rpm. The acetonitrile layer was passed through a 0.22-µm-pore-size filter for processing by HPLC.

An autosampler with an electric sample valve, a variable wavelength diode array detector, and a quaternary pump were part of the HPLC system (Agilent HPLC 1260 infinite series; Agilent Technologies, Santa Clara, CA). A 150 mm  $\times$  4.6 mm  $\times$  5 m octadecyl-silica analytical column was used in the HPLC system. The injection volume was 20 µl, the detection wavelengths were 210 and 220 nm, and the flow rates of the mobile phases (acetonitrile 90% + water 10% and acetonitrile 65% + water 35%) were 0.8 and 1 ml/min for flubendiamide and flonicamid, respectively. The retention times for flubendiamide and flonicamid were 4.6 and 4.7 min, respectively.

**Statistical analysis of residues.** The first-order kinetic model was used to describe how flonicamid and flubendiamide residues dissipated in tomato tissues and soil, as calculated with the formula of Hoskins et al. (1961):  $C_t = C_0 e^{-kt}$ , where  $C_0$  is the initial concentration (mg/kg),  $C_t$  is the residue concentration (mg/kg) at time t (day) after the pesticide treatment, and k is the degradation rate constant (per day). The half-life ( $t_{\chi_2}$ ) used to assess the dissipation rate was determined using the formula of Saber et al. (2016),  $t_{\chi_2} = \ln 2/k$ .

**Dietary exposure risk assessment.** Formulae used to calculate the risk quotient (RQ) for the long-term intake risk and the national estimated daily intake (NEDI, mg/kg/day) were those of Wang et al. (2016) and Qian et al. (2017), where NEDI = STMRi  $\times$  Fi/bw, and RQ = NEDI/ADI. In these equations, STMRi is the median residue data from controlled trials (in our case, we used the mean because we had three replicates), Fi is food intake (kg/day), bw is body weight (kg), and ADI is the acceptable daily intake. In Egypt, the average body weight of an adult is 60 kg, which was used in our calculations. RQ values <1 are regarded by consumers as acceptable risks, whereas RQ values >1 are regarded as unacceptable risks (Oliva et al. 2017, Zhang et al. 2021).

**Method validation.** The guidelines for the validation of analytical methods and quality control procedures for pesticide residues in food and feed were published by the European Commission (2019). To determine the viability of the suggested procedure, linearity, matrix effect (ME), accuracy, precision, and limit of quantification (LOQ) were all evaluated. Three concentrations of flubendiamide and flonicamid standard solutions were applied to blank samples of the tomato fruits and soil to assess the method accuracy and precision. Each treatment was replicated five times. For additional concentrations of 0.1, 0.5, and 1 mg/kg, the recovery rate and relative standard deviation (RSD) were calculated to determine the most effective combination of purifying agents. The limit of detection and the LOQ were used to test the sensitivity of the method. The lowest spiked concentration quantification was used to define the LOQ of the suggested approach.

Because coextracts produced during pretreatment affect the accuracy of the results, the ME was calculated according to the formula of Hoff et al. (2015):  $ME\% = [(k \text{ matrix} - k \text{ solvent})/k \text{ solvent}] \times 100$ , where k solvent is the slope of the solvent calibration curve and k matrix is the slope of the matrix calibration curve. An ME value of >10% indicates a definite matrix strengthening or weakening effect.

#### Results

Effectiveness test. After the application of all four the tested insecticides, B. tabaci nymph infestation decreased significantly during both the 2022 season (F = 43; df = 28; P = 0.0001) and the 2023 season (F = 48.57; df = 28; P = 0.0001) (Fig. 1). The application of flubendiamide and flonicamid resulted in lower infestation than did application of acetamiprid and azadirachtin in both seasons (Tables 2, 3). One day after the application of azadirachtin, acetamiprid, flubendiamide, and flonicamid during the 2022 season, the reductions in *B. tabaci* nymph infestation were 18.3, 67.0, 81.5, and 86.9%, respectively, whereas 3 d after application reductions were 35.98, 72.5, 86.47, and 94.49%, respectively (Table 4; Fig. 2). Residual effects were noted 5-21 d after application of the four insecticides, with mean reductions of 90.95, 85.82, 60.11, and 43.91%, respectively. A similar trend was observed in 2023 when flubendiamide and flonicamid were more effective than acetamiprid and azadirachtin (Table 5; Fig. 2). The reductions in B. tabaci infestation 1 d after of application of flubendiamide and flonicamid were 78.61 and 84.14%, respectively, and reached 89.28 and 92.71%, respectively, 3 d after application. Residual effects of flubendiamide and flonicamid resulted in mean reductions of 87.75 and 90.37%, respectively.

Effect of flubendiamide and flonicamid on detoxifying enzymes. The activities of  $\alpha$ -esterase, cytochrome P-450, and GST were determined in *B. tabaci* adults at 1, 7, 15, and 21 d following field application of flubendiamide and flonicamid. Flonicamid increased the  $\alpha$ -esterase activity of *B. tabaci* at 7, 15, and 21 d after application by 1.36, 1.97 and 1.76 times, respectively (Table 6; Fig. 3), whereas flubendiamide increased  $\alpha$ -esterase activity only 7 and 21 d after application by 1.56 and 1.68 times, respectively. In contrast, both insecticides did not significantly reduce the cytochrome P-450 activity (0.0154 ± 0.0019 and 0.0153 ± 0.004, respectively) at 21 d after application compared with the control (0.0184 ± 0.0016). In addition, flonicamid and flubendiamide did not significantly decrease the activity of GST at 7 d (8.7 ± 1.28 and 10.11 ± 4.61, respectively) and 21 d (6.58 ± 0.29 and 9.1 ± 2.46, respectively) after application compared with the control (Table 6; Fig. 3).

**Method validation.** ME, linearity, LOQ, accuracy, and precision were among the validation parameters assessed for the analytical method. The correlation coefficient ( $R^2$ ) of the flubendiamide or flonicamid calibration curve with the matrix calibration curve was used to measure the linearity. The linear range complies with the requirement for  $R^2$  (0.99). For tomato fruits, the matrix standard curve and the standard curves of the flubendiamide and flonicamid solutions yielded  $R^2$  values of >0.98, indicating a strong linear relationship. According to Matuszewski et al. (2003), ME describes how matrices other than the tested compound might affect the target's response value. In cherry tomato fruits, the MEs of flonicamid and flubendiamide were  $\geq$ 0.98. No measurable ME was present. As a result, the effect of other matrices on the target compound's response value was minimal. Flubendiamide and flonicamid in cherry tomatoes could be quantified only to a maximum of 0.01 mg/kg.

The recovery test and RSD were used to confirm the procedure's accuracy and precision. The recovery rates of flubendiamide and flonicamid from tomato obtained utilizing the ideal purifier combination are shown in Table 7. The recovery





Fig. 1. Mean ( $\pm$ SD) number of *B. tabaci* nymphs on cherry tomato plants after field application of four insecticides during the 2022 and 2023 seasons.

rates and the matching RSD of flubendiamide and flonicamid are shown for the three spiking levels in fruits and soil. Thus, by using the suggested procedure, high recoveries of flubendiamide and flonicamid at the three spiking levels were obtained within the acceptance criterion of 80–110%, with a precision RSD of  $\geq$ 20% (European Commission 2019).

four	
n of	
applicatio	
field	
after	
and	
before	
plants	
tomato	
cherry	
s on	
nymph	season.*
tabaci	owing
of B.	2 gr
umber o	the 202
SD) n	ides ir
+  u	ctic
Mea	inse
Table 2.	

	Before				After Application			
Treatment	Application	Day 1	Day 3	Day 5	Day 7	Day 10	Day 15	Day 21
Control	101.3 ± 5.11a	118 ± 5.14a	122.8 ± 1.92a	133.3 ± 4.91a	148 ± 10.79a	155.5 ± 3.64a	169 ± 10.29a	186 ± 4.47a
Azadirachtin	103 ± 3.08a	$98.8 \pm 4.96b$	$80.5 \pm 4.5b$	$68 \pm 9.08b$	$65.8 \pm 7.66b$	$85 \pm 4.3b$	$108.5 \pm 13.12b$	$135.5 \pm 6.1b$
Acetamiprid	97.5 ± 2.17ab	$37.5 \pm 2.69c$	$32.5 \pm \mathbf{4.09c}$	$30 \pm 6.67c$	$51.3 \pm 5.44b$	$63.8 \pm 3.34c$	$72.5 \pm 7.43c$	$95 \pm 6.59c$
Flubendiamide	98 ± 3.93ab	$15 \pm 1.22d$	$6.25 \pm 2.94e$	$3.8 \pm 3.26d$	$6.3 \pm 2.77c$	$12.5 \pm 2.17d$	$21.3 \pm 4.43d$	$30 \pm 7.71d$
Flonicamid	$91.5 \pm 2.29b$	$20 \pm 2.44d$	$15 \pm 3.16d$	$10 \pm 2.23d$	$11.3 \pm 3.96c$	$17.5 \pm 6.1d$	$26.3 \pm 6.6d$	$37.5 \pm \mathbf{5.31d}$
* Mithin o column	a moone with different	tottore are cignific	The second second second second second					

Within a column, means with different letters are significantly different (Tukey's HSD, P = 0.05).

Table 3. Mean ( $\pm$ SD) number of *B. tabaci* nymphs on cherry tomato plants before and after field application of four insecticides in 2023 growing season.\*

	Before				After Application	_		
Treatment	Application	Day 1	Day 3	Day 5	Day 7	Day 10	Day 15	Day 21
Control	92 ± 3.16bc	98.8 ± 3.11a	105.8 ± 3.76a	121 ± 8.24a	129.3 ± 7.11a	146 ± 7.64a	155.5 ± 10.64a	188.5 ± 5.72a
Azadirachtin	98 ± 3.16ab	96 ± 2.54a	$77.8 \pm 2.86b$	$70.8 \pm 5.93b$	$60.5 \pm 5.54b$	98 ± 3.8b	$112.5 \pm 8.64b$	$145 \pm 1.58b$
Acetamiprid	$88.3 \pm 2.86c$	$35 \pm 2.54b$	$31.8 \pm 7.49c$	$46.3 \pm 4.14c$	$59.3 \pm 5.35b$	$69.3 \pm 5.11c$	$76.3 \pm 13.64c$	$102.5 \pm 5.02c$
Flubendiamide	$95.5 \pm 2.95 bc$	$16.3 \pm 2.27c$	$8 \pm 2.73d$	$5.3 \pm 1.47d$	$6.8\pm\mathbf{2.86c}$	$13.3 \pm 2.38d$	$19.3 \pm 6.17d$	33.8 ± 8.58d
Flonicamid	$103.5 \pm 2.29a$	$23.8 \pm 4.49c$	$12.8 \pm 4.14d$	$\textbf{9.3} \pm \textbf{4.81d}$	$11.5 \pm 3.77c$	$21.3 \pm 4.81d$	$25.3 \pm 6.33d$	$41.3 \pm 6.86d$
* 146461	ter en e Affie - differ e e e e e e e	and and and another the						

Within a column, means with different letters are significantly different (Tukey's HSD, P = 0.05).

fou	r insecticides	in the 2022 g	rowing seaso	'n.*				
Insecticide	Day 1	Day 3	Day 5	Day 7	Day 10	Day 15	Day 21	Average Residual Effect**
Azadirachtin	18.3 ± 1.41a	36 ± 2.2a	50.2 ± 3.62c	56.7 ± 2.89b	46.7 ± 5.68c	37.1 ± 5.19c	29 ± 3.09c	43.9 ± 10.67
Acetamiprid	$67 \pm 2.12b$	$72.5 \pm 3.2b$	$76.6 \pm 3.44b$	$64.1 \pm 2.98b$	$57.4 \pm 3.72b$	$55.5 \pm 3.73b$	$47 \pm 4.31b$	$60.1 \pm 10.55$
Flubendiamide	$81.5 \pm 2.66c$	$86.5 \pm 2.33c$	91.7 ± 2.4a	89.9 ± 4.74a	87.3 ± 4.26a	82.6 ± 3.89a	77.7 ± 2.65a	$85.8 \pm 6.3$
Flonicamid	$86.9 \pm 1.58d$	$94.5 \pm 2.63d$	97.1 ± 2.3a	95.6 ± 1.89a	91.7 ± 3.04a	87 ± 2.55a	83.4 ± 3.82a	$90.9 \pm 5.88$
* Within a column	means with differe	ant lattars ara signif	inanthy different (Tu	kev's HSD $P = 0.0$	75)			

.(cn.n = Within a column, means with different letters are significantly different (1 ukey's HSU, F

\*\* Average residual effect was calculated at 5–21 d after application.

Table 4. Mean (±SD) percent reduction in *B. tabaci* nymph infestation on cherry tomato plants after field application of



Fig. 2. Step slope chart representing the mean ( $\pm$ SD) percent reductions in *B. tabaci* nymphs after field application of four insecticides during the 2022 and 2023 seasons.

Dissipation and terminal residue of flubendiamide and flonicamid in fruits, leaves, and soil. The dissipation of flubendiamide and flonicamid in cherry tomato fruits and leaves and in soil was determined. The dissipation kinetics and half-life are displayed in Tables 8 and 9. In fruits, leaves, and soil, flubendiamide and flonicamid had half-lives of 4.25-2.60 d. The final residual amounts of flubendiamide and flonicamid in fruits, leaves, and soil are displayed in Tables 8 and 9. Flubendiamide had an initial residue of 3.80, 6.33, and 1.30 mg/kg in fruits, leaves, and soil, respectively (Table 8). After 1 d, the residue concentration dropped to 2.95, 5.84, and 0.99 mg/kg, respectively, with losses of 22.36, 7.74, and 23.84%, respectively. At 15 d following treatment, the concentrations in fruits, leaves, and soil decreased to 0.05, 0.1, and 0.07 mg/kg, respectively, with losses of 98.68, 98.42, and 94.61%, respectively. At 21 d after treatment, the residues in fruits and soil were below the detection limits. Thus, if the maximum residue level (MRL) for tomato were 2 mg/kg, the recommended preharvest interval (PHI) would be 3 d if the specified dosage and safety interval recommendations were followed. The residues of flonicamid in fruits, leaves, and soil were 4.08, 7.51, and 3.72 mg/kg, respectively, 2 h after treatment (Table 9). At 1 d, the residues dropped to 3.36, 5.57, and 2.17, respectively. At 3 d after application, rapid dissipation was noted in fruits, leaves, and soil, with flonicamid residues declining to 2.64, 4.33, and 1.58 mg/kg, respectively, and losses of 35.29, 42.34, and 57.52%, respectively. After 15 d, the residues fell to 0.05 and 0.16 mg/kg and not detectable, with 98.77, 97.86, and 100% loss, respectively. Flonicamid residues were no longer identifiable at 21 d following application. Accordingly, the PHI recommendation is 7 d when using the specified dosage and based on an MRL of 0.5 mg/kg for cherry tomato (Table 9).

for	ir insecticide	s in the 2022 (	growing seas	on.*				
Insecticide	Day 1	Day 3	Day 5	Day 7	Day 10	Day 15	Day 21	Average Residual Effect**
Azadirachtin	12.7 ± 2.11c	31 ± 1.87c	45.1 ± 4.58c	56.6 ± 5.07b	37 ± 3.16c	$32.4 \pm 3.18c$	27.8 ± 1.9c	<b>39.8</b> ± <b>10.83</b>
Acetamiprid	$62.3 \pm 3.29b$	$68.7 \pm \mathbf{6.18b}$	$60.2 \pm 2.84b$	$52.2 \pm 3.16b$	$50.6 \pm 3.17b$	$48.9 \pm 5.37b$	$43.3 \pm 2.43b$	$51.02 \pm 6.51$
Flubendiamide	78.6 ± 4.04a	89.3 ± 3.23a	93.5 ± 3.67a	92.1 ± 3.05a	87.3 ± 2.79a	85.4 ± 3.37a	80.6 ± 3.14a	$87.8 \pm 5.67$
Flonicamid	84.1 ± 3.02a	92.7 ± 2.64a	95.8 ± 1.79a	95 ± 2.25a	91 ± 2.24a	88.1 ± 4.24a	82 ± 4.3a	$90.4 \pm 5.94$
* Within a column	meane with differ	rant lattare are cian	ificantly different (T	- I USH Street	0.05\			

= u.uoj. Within a column, means with different letters are significantly different (Lukey's HSD, P

\*\* Average residual effect was calculated at 5–21 d after application.

Table 5. Mean ( $\pm$ SD) percent reduction in *B. tabaci* nymph infestation on cherry tomato plants after field application of

Table 6. Mean ( $\pm$ SD) enzymatic activity of  $\alpha$ -esterase, cytochrome P-450, and GST in *B. tabaci* adults at 1, 7, 15, and 21 d following field application of flubendiamide and flonicamid.\*

			Enzyme Activity (I	Mmole/mg protein)	
Enzyme	Treatment	Day 1	Day 7	Day 15	Day 21
α-esterase	Control	23.94 ± 9.91a	18.21 ± 6.008a	11.35 ± 0.67a	7.32 ± 1.42a
	Flubendiamide	20.15 ± 8.05a	28.57 ± 19.1a	8.9 ± 0.9a	12.31 ± 4.94a
	Flonicamid	17.72 ± 5.11a	24.82 ± 8.22a	22.42 ± 8.04a	12.95 ± 2.92a
P-450	Control	0.019 ± 0.0054a	$0.0202\pm0.0032a$	0.0137 ± 0.0027a	$0.0184 \pm 0.0016a$
	Flubendiamide	0.0209 ± 0.0043a	0.0268 ± 0.0161a	0.0141 ± 0.0014a	0.0153 ± 0.004a
	Flonicamid	$0.0202 \pm 0.0007a$	$0.0182 \pm 0.0027a$	0.0194 ± 0.0067a	0.0154 ± 0.0019a
GST	Control	8.76 ± 1.54a	10.27 ± 1.6a	9.29 ± 0.73a	9.58 ± 0.59a
	Flubendiamide	9.67 ± 1.06a	10.11 ± 4.61a	6.72 ± 0.9a	9.1 ± 2.46a
	Flonicamid	10.08 ± 0.2a	8.7 ± 1.28a	10 ± 3.71a	6.58 ± 0.29a

\* Within a column, means with different letters are significantly different (Tukey's HSD, P = 0.05).

**Risk assessment.** The RQ is of crucial significance to human health safety; it is calculated by dividing the NEDI by the ADI. An RQ value of <1 indicates low potential risk to consumers, and a value of >1 means high potential risk. For long-term consumption, flubendiamide and flonicamid had a high risk for the first 5 d of use on tomatoes treated with the recommended concentrations. According to our findings, the RQ was >1 for 0, 1, 3, and 5 d following application. However, for days 7, 10, and 15, the RQ values were <1, which indicates a low risk for human health (Table 10).

# Discussion

Chemical control is considered one of the preferred methods for quick relief from insect pests. However, pesticide application can yield questionable outcomes



Fig. 3. Radar chart representing the activities of  $\alpha$ -esterase, cytochrome P-450, and GST in *B. tabaci* adults at 1, 7, 15, and 21 d after field application of flubendiamide and flonicamid.

Table 7. Meai	n (±SD) percentage	recovei	y of flubendiamide	and flor	nicamid residues fro	m cherr	y tomato fruits and	soil.*
		Flubend	diamide			Flonic	amid	
Spiked level	Tomato fruit	ő	Soil		Tomato fruits	6	Soil	
(b/brl)	Recovery ± RSD	RSD,	Recovery ± RSD	RSD,	Recovery ± RSD	RSD,	Recovery ± RSD	RSD,
0.1	$92.76 \pm 0.40$	0.46	$85.36 \pm 3.3$	3.86	$95.97 \pm 1.50$	1.58	$81.32 \pm 2.10$	2.52
0.5	$95.10 \pm 0.67$	0.70	$85.81 \pm 1.03$	1.20	$97.29 \pm 2.58$	2.65	$84.02 \pm 1.42$	1.68
-	$98.24 \pm 1.01$	1.02	$95.71 \pm 1.53$	1.60	$106.0 \pm 1.67$	1.57	$89.72 \pm 2.65$	2.95

\* RSD = relative standard deviation; RSD<sub>r</sub> = recovery repeatability (precision).

Table 8. Dissipation kinetics of flubendiamide residues (mean  $\pm$  SD) in cherry tomato fruits and leaves and in soil under onen field conditions.

		2							
	To	mato Fri	uits	Ton	nato Lea	ves		Soil	
Day After Application	Residue (mg/kg)	% Loss	% Persistence	Residue (mg/kg)	% Loss	% Persistence	Residue (mg/kg)	% Loss	% Persistence
0	$3.80 \pm 1.25$	00.0	100.0	$6.33 \pm 1.05$	0.00	100.0	$1.30 \pm 0.52$	0.00	100.0
+	$2.95 \pm 1.04$	22.36	77.64	$5.84 \pm 1.09$	7.74	92.26	$0.99 \pm 0.63$	23.84	76.16
с	$1.82 \pm 0.95$	52.10	47.90	$3.72 \pm 0.92$	41.23	58.77	$0.77 \pm 0.19$	40.76	59.24
5	$0.98 \pm 2.01$	74.21	25.79	$\textbf{2.90} \pm \textbf{0.48}$	54.18	45.82	$0.39 \pm 0.47$	70.00	30.00
7	$0.45 \pm 1.44$	88.15	11.85	$1.25 \pm 0.94$	80.25	19.75	$0.24 \pm 1.37$	81.53	18.47
10	$0.27 \pm 0.98$	92.89	7.11	$0.57~\pm~1.32$	90.99	9.01	0.11 ± 1.26	91.53	8.47
15	$0.05 \pm 2.27$	98.68	1.32	$0.10 \pm 1.29$	98.42	1.58	$0.07 \pm 1.50$	94.61	5.39
21	ND*			$\textbf{0.06} \pm \textbf{2.00}$	99.05	0.95	QN		
$t_{1_{2_{2}}}$ (days)		3.13			3.63			3.68	
MRL** (mg/kg)		0							
PHI† (day)		ი							
* ND - not detectable									

\*\* MRL = maximum residue level.

† PHI = preharvest interval.

open	
under	
soil	
and	
leaves,	
fruits,	
tomato	
cherry	
<u>ב</u> .	
SD)	
+1	
(mean	
residues	
flonicamid	
of	
kinetics	ons.
Dissipation	field conditi
9.	
Table	

	To	mato fru	lits	Tor	nato lea	ves		Soil	
Day After Application	Residue (mg/kg)	% Loss	% Persistence	Residue (mg/kg)	% Loss	% Persistence	Residue (mg/kg)	% Loss	% Persistence
0	$4.08 \pm 0.98$	00.0	100.0	7.51 ± 0.82	00.0	100.0	$3.72 \pm 052$	0.00	100.0
-	$3.36 \pm 0.47$	17.64	82.36	$5.57 \pm 0.48$	25.83	74.17	$2.17 \pm 0.93$	41.66	58.34
ო	$2.64 \pm 1.20$	35.29	64.71	$4.33 \pm 0.55$	42.34	57.66	$1.58 \pm 0.22$	57.52	42.48
5	$1.09 \pm 0.85$	73.28	26.72	$\textbf{2.22} \pm \textbf{0.21}$	70.43	29.57	$\textbf{0.84}\pm\textbf{0.17}$	77.41	22.59
7	$0.42 \pm 0.56$	89.7	10.30	$\textbf{1.31}\pm\textbf{0.63}$	82.55	17.45	$0.31 \pm 0.82$	91.66	8.34
10	$0.13 \pm 1.83$	96.81	3.19	$0.24\pm0.85$	96.80	3.20	$\textbf{0.11}\pm\textbf{0.55}$	97.04	2.96
15	$0.05 \pm 1.22$	98.77	1.23	$0.16 \pm 0.99$	97.86	2.14	ND*		
21	QN			$0.04 \pm 1.36$	99.46	0.54	ND		
$t_{1_{2_{2}}}$ (days)		4.25			3.54			2.60	
MRL** (mg/kg)		0.5							
PHI† (day)		7							
* ND = not detectable ** MRL = maximum re	esidue level.								

304

J. Entomol. Sci. Vol. 59, No. 3 (2024)

Downloaded from https://prime-pdf-watermark.prime-prod.pubfactory.com/ at 2025-07-01 via free access

 $\uparrow$  PHI = preharvest interval.

	Flub	endiamide	)	Fle	onicamid	
Day After Application	Residue (mg/kg)	NEDI	RQ	Residue (mg/kg)	NEDI	RQ
0	3.8	0.82	8.17	4.08	0.88	8.77
1	2.95	0.63	6.34	3.36	0.72	7.22
3	1.82	0.39	3.91	2.64	0.57	5.68
5	0.98	0.21	2.11	1.09	0.23	2.34
7	0.45	0.10	0.97	0.42	0.09	0.90
10	0.27	0.06	0.58	0.13	0.03	0.28
15	0.05	0.01	0.11	0.05	0.01	0.11
21	ND*	ND	ND	ND	ND	ND

Table 10. National estimated daily intake (NEDI) and risk quotient (RQ) of flubendiamide and flonicamid in cherry tomato fruits.

\* ND = not detectable.

due to several factors (Moustafa et al. 2018). Therefore, the residue dissipation pattern, half-life, and PHI must be evaluated for each pesticide. The purpose of the current study was to study the effectiveness of azadirachtin, acetamiprid, flubendiamide, and flonicamid against *B. tabaci* infestation and to determine the biochemical impact and persistence of flubendiamide and flonicamid residues in cherry tomato fruits and leaves and in soil samples.

Our results clearly indicate that the initial kill and residual effect on the whitefly (B. tabaci) nymphs in the 2022 season were the same as those obtained in 2023. Flonicamid and flubendiamide resulted in the highest reduction in *B. tabaci* infestation, followed by acetamiprid and azadirachtin. These results agree with those of Assadi et al. (2022), who reported significant insecticidal activity of flonicamid against eggs, nymphs, and adults of *B. tabaci*, indicating that this pesticide could be used effectively for B. tabaci management (Huded et al. 2019, Kodandaram et al. 2017, Roditakis et al. 2014, Sadhana et al. 2021). Flubendiamide achieved adequate control of whitefly nymphs (Dake and Bhamare 2019), but acetamiprid effectiveness was limited due to the resistance of field populations of piercingsucking insects to acetamiprid and other insecticides (Koo et al. 2014, Ullah et al. 2020, Wang et al. 2007). Azadirachtin has was less effective against B. tabaci, as occurs with other neem (Azadirachta indica Juss) products. Azadirachtin contains triterpenoids, and these active ingredients undergo rapid degradation by ultraviolet radiation when used as a foliar application (Barnaby et al. 1989, Barrek et al. 2004, Johnson et al. 2003). Mortality of B. tabaci under field conditions were 55.4-67.0% with azadirachtin (Prijović et al. 2012).

Detoxification enzymes are the key factors in the metabolism of toxic compounds in insects (Moustafa et al. 2023d, Prasannakumar et al. 2023). Therefore, insect resistance is usually associated with increased activity of these enzymes (Fouad et al. 2022). In the current study, the enzymatic activity of esterase did not increase significantly in the nymphs after flubendiamide and flonicamid application. Thus, esterase enzyme activity could serve as an indicator of whitefly adaptation to insecticides. In contrast, both insecticides did not decrease the activities of cytochrome P-450 and GST, which may be a major cause of whitefly nymph mortality.

Pesticide residues in plants are significantly influenced by a number of essential elements, including the photolysis and volatilization of pesticides brought on by high temperatures and light, rain, and the physical and chemical properties of pesticides (Subirats et al. 2005). With regards to residues, our results for both flubendiamide and flonicamid in cherry tomato fruits and leaves and in soil samples agreed with those of Kelageri et al. (2017), who reported 1.23 mg/kg in tomato samples taken from polyhouses; this residue diminished to below detectable limits by day 10, with a half-life of 6.18 d. Deposits of 0.90 mg/kg in open fields diffused to below detectable limits by day 7, with a half-life of 6.07 d, indicating that polyhouse dissipation was slower than that in the open fields for a variety of reasons.

According to the Organization for Economic Cooperation and Development calculator and an assessment of the chronic hazard exposure factoring in average body weight, national tomato consumption per capita, and the ADI of flubendiamide, MRLs of 3 mg/kg for polyhouse tomatoes and 2 mg/kg for open field tomatoes are advised. A 4% acetic acid solution (61.63%) was the least effective decontamination method (17.71%) for removing flubendiamide residues from cherry tomatoes, whereas a vegetable wash was the most effective procedure for removing flubendiamide residues (65.39%) and can be recommended as a risk mitigation method for food safety.

Our findings were also consistent with those of Wang et al. (2018), who investigated the persistence of flonicamid and its metabolites in soil and cabbage (*Brassica* sp.) after harvest and their dissipation behavior. The half-lives of flonicamid alone and total residues (the sum of flonicamid and its metabolites) were 1.49– 4.59 and 1.97–4.99 d, respectively, in cabbage plants and 2.12–7.97 and 2.04– 7.62 d, respectively, in soil. Flonicamid decomposed quickly. When 50% flonicamid WG was sprayed once or twice at the recommended dose and at 1.5 times the recommended dose, the highest residues of total flonicamid in cabbage and soil at various PHIs (3, 7, and 14 d) were 0.070 and 0.054 mg/kg, respectively.

According to consumption data from China, flonicamid's RQ was <16.84%, which suggests that using it is safe for people. These findings could assist the Chinese government in determining the MRL for flonicamid in cabbage and could provide guidance for the safe and responsible use of flonicamid in agriculture. Li et al. (2021) found that total and chlorfluazuron RQs were both <1. This finding suggests that there is very little dietary danger from chlorfluazuron in tea. However, compared with chlorfluazuron, total flonicamid ingestion carries a threefold greater risk. Consuming tea containing flonicamid and its metabolites poses a risk to human health. In contrast, Kelageri et al. (2017) found that flubendiamide in tomatoes has low risk, and the total dry matter intake was less than the ADI in open fields and polyhouses. Jankowska et al. (2022) investigated the long-term risk of pesticide residues on some vegetables. The RQ was >1 for flonicamid in Brussels sprouts (Brassica sp.), which indicated negative health effects on humans. This finding is not consistent with those of our study, where the RQ was high for only 5 d and then decreased. Evaluation of the dissipation and health risk for some insecticides in tomatoes under field and polyhouse conditions revealed that no health risk for consumers for the tested pesticides, including flubendiamide (Singh et al. 2023). These results support the accuracy and precision of the proposed procedure, affirming its suitability for the quantitative analysis of flubendiamide and flonicamid in cherry tomato samples.

In conclusion, flubendiamide and flonicamid could be used as effective insecticides in the whitefly management program in Egypt, and this program should include various groups of insecticides. Flubendiamide and flonicamid concentrations in diverse matrices were determined quickly and effectively in our study with a QuEChERS technique. For flubendiamide and flonicamid, the PHI were 3 and 7 d, respectively. Estimations of food exposure become are crucial when investigating pesticide residues.

#### Acknowledgment

Many thanks are due to Dr. Hamed Khairullah Said (Associate Professor, Department of Economic Entomology and Pesticides, Faculty of Agriculture, Cairo University, Egypt) for revising and editing the manuscript for grammatical correctness and coherence.

# **References Cited**

- Anastassiades, M., S.J. Lehotay, D. Tajnbaher and J.F. Schenck. 2003. Fast and easy multiresidues method employing acetonitrile extraction/partitioning and "dispersive solidphase extraction" for the determination of pesticide residues in produce. J. AOAC. Int. 86: 412–430. https://doi.org/10.1093/jaoac/86.2.412
- Assadi, B.H., S. Chouikhi and M.S. Belkadhi. 2022. Effectiveness of flonicamid 50WG against *Bemisia tabaci* (Genn.) under greenhouse conditions in Tunisia. J. Oasis Agric. Sustain. Dev. 4(2): 214–217. https://doi.org/10.56027/JOASD.spiss282022
- Barnby, M.A., R.B. Yamasaki and J.A. Kloske. 1989. Biological activity of azadirachtin, three derivatives, and other ultraviolet radiation degradation products against tobacco budworm (Lepidoptera: Noctuidae) larvae. J. Econ. Entomol. 82: 58–63.
- Barrek, S., O. Paisse and G.L. Marie-Florence. 2004. Analysis of neem oils by LC-MS and degradation kinetics of azadirachtin-A in a controlled environment. Anal. Bioanal. Chem. 378: 753–763. https://doi.org/10.1007/s00216-003-2377-0
- **Casida, J.E. 2015.** Golden age of RyR and GABA-R diamide and isoxazoline insecticides: Common genesis, serendipity, surprises, selectivity, and safety. Chem. Res. Toxicol. 28: 560–566. https://doi.org/10.1021/tx500520w
- Cordova, D., E.A. Benner, M.D. Sacher, J.J. Rauh, J.S. Sopa, G.P. Lahm, T.P. Selby, T.M. Stevenson, L. Flexner, S. Gutteridge, D.F. Rhoades, L. Wu, R.M. Smith and Y. Tao. 2006. Anthranilic diamides: A new class of insecticides with a novel mode of action, ryanodine receptor activation. Pestic. Biochem. Physiol. 84: 196–214. https://doi.org/10. 1016/j.pestbp.2005.07.005
- Dake, R.B. and V.K. Bhamare. 2019. Bio-efficacy, persistence and residual toxicity of different insecticides against whitefly (*Bemisia tabaci* (Gennadius)) on sunflower. J. Entomol. Zool. Stud. 7: 1499–1504.
- El-Hefny, D.E., H. Hunter and R.M.A. Helmy. 2021. Validation using QuEChERS method, risk assessment and preharvest intermission using GC-MS for determination of azoxystrobin in tomato and cucumber. Egypt J. Chem. 64: 7421–7429. https://doi.org/10. 21608/EJCHEM.2021.76725.3756
- **European Commission. 2019.** Analytical quality control and method validation procedures for pesticide residues analysis in food and feed. Document No SANTE/12682/2019.
- Fouad, E.A., F.S. Ahmed and M.A.M. Moustafa. 2022. Monitoring and biochemical impact of insecticides resistance on field populations of *Spodoptera littoralis* (Boids.) (Lepidoptera: Noctuidae) in Egypt. Polish J. Entomol. 91: 109–118.

- Ghosal, A. and M. Chatterjee. 2018. Insecticide induced resurgence study of whitefly in cotton and tomato. Univ. Sin. J. Anim. Sci. 2: 1–6. https://doi.org/10.13140/RG.2.2.30985.08805
- Habig, W.H., M.J. Pabst and W.B. Jakoby. 1974. Glutathione S-transferases: The first enzymatic step in mercapturic acid formation. J. Biol. Chem. 249: 7130–7139.
- Hansen, L.G. and E. Hodgson. 1971. Biochemical characteristics of insect microsomes: Nand O-demethylation. Biochem. Pharmacol. 20: 1569–1578. https://doi.org/10.1016/ 0006-2952(71)90285-1
- Henderson, C.F. and E.W. Tilton. 1955. Tests with acaricides against the brown wheat mite. J. Econ. Entomol. 48: 157–161. https://doi.org/10.1093/jee/48.2.157
- Hoff, R.B., G.L.F. Rübensam Jank, M.C.R. Barreto, T.M. Peralba, M.S. Pizzolato and D. Barceló Díaz-Cruz. 2015. Analytical quality assurance in veterinary drug residue analysis methods: Matrix effects determination and monitoring for sulfonamides analysis. Talanta 132: 443–450. https://doi.org/10.1016/j.talanta.2014.08.046
- Hoskins, W.M. 1961. Mathematical treatment of loss of pesticide residues. Plant Prot. Bull. 9: 163–168.
- Huded, P.B., V.N. Ghante and V. Kulkarni. 2019. Chemical management of whiteflies in sunflower. Int. J. Curr. Microbiol. Appl. Sci. 8: 2885–2890. https://doi.org/10.20546/ ijcmas.2019.804.337
- Jankowska, M., I. Hrynko and B. Łozowicka. 2022. Human health risk assessment of pesticide residues in fruit, vegetable and cereal samples from Poland—A 5-year survey. J. Plant Prot. Res. 62: 385–392. https://doi.org/10.24425/jppr.2022.143231
- Johnson, S., P. Dureja and S. Dhingra. 2003. Photostablizers for azadirachtin-A (a neembased pesticide). J. Environ. Sci. Health B 38: 451–462. https://doi.org/10.1081/PFC-120021665
- Kadala, A., M. Charreton and C. Collet. 2020. Flubendiamide, the first phthalic acid diamide insecticide, impairs neuronal calcium signalling in the honey bee's antennae. J. Insect Physiol. 125: 104086. https://doi.org/10.1016/j.jinsphys.2020.104086
- Kandil, M.A., M.A.M. Moustafa, M.A. Saleh and I.R. Ateya. 2023. Dissipation kinetics and degradation products of cyantraniliprole in tomato plants and soil in the open field. Egyptian J. Chem. 66: 483–493. https://doi.org/10.21608/EJCHEM.2023.195133.7625
- Kelageri, S.S., C.S. Rao, V.S. Bhushan, P.N. Reddy, S.S. Rani, M. Aruna, A.H. Reddy, D. Ravindranath, B. Ramesh and M. Hymavathy. 2017. Flubendiamide: Residues and risk assessment in tomato Solanum lycopersicum. Int. J. Agric. Environ. Biotechnol. 10: 651–667. https://doi.org/10.5958/2230-732X.2017.00081.X
- Kodandaram, M.H., Y.B. Kumar, K. Banerjee, S. Hingmire, A.B. Rai and B. Singh. 2017. Field bioefficacy, phytotoxicity and residue dynamics of the insecticide flonicamid (50 WG) in okra [Abelmoschus esculenta (L) Moench]. Crop Prot. 94: 13–19. https://doi.org/ 10.1016/j.cropro.2016.12.003
- Koo, H.N., J.J. An, S.E. Park, J.I. Kim and G.H. Kim. 2014. Regional susceptibilities to 12 insecticides of melon and cotton aphid, *Aphis gossypii* (Hemiptera: Aphididae) and a point mutation associated with imidacloprid resistance. Crop Prot. 55: 91–97.
- Kumar, V., V. Jindal, S.K. Kataria and M. Pathania. 2019. Activity of novel insecticides against different life stages of whitefly (*Bemisia tabaci*). Indian J. Agric. Sci. 89: 1599– 1603. https://doi.org/10.56093/ijas.v89i10.94585
- Li, H., Q. Zhong, X. Wang, F. Luo, L. Zhou, H. Sun, M. Yang, Z. Lou, Z. Chen and X. Zhang. 2021. The degradation and metabolism of chlorfluazuron and flonicamid in tea: A risk assessment from tea garden to cup. Sci. Total Environ. 754: 142070. https://doi.org/ 10.1016/j.scitotenv.2020.142070
- Li, W., Y. Zhang, H.R. Jia, W.W. Zhou, B.T. Li and H.J. Huang. 2019. Residue analysis of tetraniliprole in rice and related environmental samples by HPLC/MS. Microchem. J. 150: 104168. https://doi.org/10.1016/j.microc.2019.104168
- Lin, T., X. Chen, L. Wang, X. Fang, M. Li, Y. Li and H. Liu. 2021. Simultaneous determination of eight new generation amide insecticide residues in complex matrix agri-food by multi-walled carbon nanotube cleanup and UPLC-MS/MS. Sci. Rep. 11: 23208. https:// doi.org/10.1038/s41598-021-02645-w

- Matuszewski, B.K., M.L. Constanzer and C.M. Chavez-Eng. 2003. Strategies for the assessment of matrix effect in quantitative bioanalytical methods based on HPLCMS/MS. Anal. Chem. 75: 3019. https://doi.org/10.1021/ac020361s
- McKenzie, C.L., V. Kumar, C.L. Palmer, R.D. Oetting and L.S. Osborne. 2014. Chemical class rotations for control of *Bemisia tabaci* (Hemiptera: Aleyrodidae) on poinsettia and their effect on cryptic species population composition. Pest Manag. Sci. 70: 1573–1587. https://doi.org/10.1002/ps.3736
- Moodley, V., A. Gubba and P.L. Mafongoya. 2019. A survey of whitefly transmitted viruses on tomato crops in South Africa. Crop Prot. 123: 21–29. https://doi.org/10.1016/j.cropro. 2019.05.018
- Morita, M., T. Ueda, T. Yoneda, T. Koyanagi and T. Haga. 2007. Flonicamid, a novel insecticide with a rapid inhibitory effect on aphid feeding. Pest Manag. Sci. 63: 969–973. https://doi.org/10.1002/ps.1423
- Morita, M., T. Ueda, T. Yoneda, T. Koyanagi, S. Murai, N. Matsuo and B. Stratman. 2000. IKI-220—A novel systemic aphicide, Pp. 59–65. *In* Brighton Crop Protection Conference—Pests and Diseases. British Crop Protection Council, Major Print Ltd., Nottingham, United Kingdom.
- Moustafa, M.A.M., A. Amer, L.A. Al-Shuraym, E.S. Ibrahim, D.E. El-Hefny, M.Z.M. Salem and S. Sayed. 2022. Efficacy of chemical and bio-pesticides on cowpea aphid, *Aphis craccivora*, and their residues on the productivity of fennel plants (*Foeniculum vulgare*). J. King Saudi Univ. Sci. 34: 101900. https://doi.org/10.1016/j.jksus.2022.101900
- Moustafa, M.A.M., D.E. El-Hefny, R.N. Abdel-kerim and M.A. Kandil. 2023a. Toxicity of insecticides against tomato leaf miner, *Tuta absoluta*, and its predators and determination of their residue dissipation in tomato fruits. J. Agric. Sci. Technol. 25: 1115–1128.
- Moustafa, M.A.M., E.A. Fouad, E. Ibrahim, A.L. Erdei, Z. Kárpáti and A. Fónagy. 2023b. The comparative toxicity, biochemical and physiological impacts of chlorantraniliprole and indoxacarb on *Mamestra brassicae* (Lepidoptera: Noctuidae). Toxics 11: 212. https://doi. org/10.3390/toxics11030212
- Moustafa, M.A.M., N.N. Hassan, N.A. Alfuhid, A. Amer and M. Awad. 2023c. Insights into the toxicity, biochemical activity and molecular docking of *Cymbopogon citratus* essential oils and citral on the cotton leafworm *Spodoptera littoralis*. J. Econ. Entomol. 116: 1185– 1195. https://doi.org/10.1093/jee/toad093
- Moustafa, M.A.M., R.I.A. Moteleb, Y.F. Ghoneim, S.S. Hafez, R.E. Ali, E.E.A. Eweis and N.N. Hassan. 2023d. Monitoring resistance and biochemical studies of three Egyptian field strains of *Spodoptera littoralis* (Lepidoptera: Noctuidae) to six insecticides. Toxics 11: 211. https://doi.org/10.3390/toxics11030211
- Moustafa, M.A.M., M.A. Saleh, I.R. Ateya and M.A. Kandil. 2018. Influence of some environmental conditions on stability and activity of *Bacillus thuringiensis* formulations against the cotton leaf worm, *Spodoptera littoralis* (Boisd.) (Lepidoptera: Noctuidae). Egypt. J. Biol. Pest Control 28: 1–7. https://doi.org/10.1186/s41938-018-0064-x
- Mugerwa, H., J. Colvin, T. Alicai, C.A. Omongo, R. Kabaalu, P. Visendi, P. Sseruwagi and S.E. Seal. 2021. Genetic diversity of whitefly (*Bemisia* spp.) on crop and uncultivated plants in Uganda: Implications for the control of this devastating pest species complex in Africa. J. Pest Sci. 94: 1307–1330. https://doi.org/10.1007/s10340-021-01355-6
- Ochilo, W.N., G.N. Nyamasyo, D. Kilalo, W. Otieno, M. Otipa, F. Chege, T. Karanja and E.K. Lingeera. 2019. Ecological limits and management practices of major arthropod pests of tomato in Kenya. J. Agric. Sci. 4: 29–42. https://doi.org/10.31248/JASP2019.124
- Oliva, J., S. Cermeno, M.A. Camara, G. Martinez and A. Barba. 2017. Disappearance of six pesticides in fresh and processed zucchini, bioavailability and health risk assessment. Food Chem. 229: 172–177. https://doi.org/10.1016/j.foodchem.2017.02.076
- Prasannakumar, N.R., N. Jyothi, S. Saroja and A.N. Lokesha. 2023. Insecticidal properties of Ocimum basilicum and Mentha piperita essential oils against South American tomato moth, Phthorimaea absoluta (Meyrick) (Lepidoptera: Gelichiidae). Pestic. Biochem. Physiol. 190: 105329. https://doi.org/10.1016/j.pestbp.2022.105329

- Prijović, M., T. Drobnjaković, D. Marčić, P. Perić and S. Petronijević. 2012. Efficacy of insecticides of natural origin in whitefly (*Trialeurodes vaporariorum*) control in tomato. Acta Hortic. 960: 359–364. https://www.researchgate.net/publication/286029175
- Qian, Y., H. Matsumoto, X. Liu, S. Li, X. Liang, Y. Liu, G. Zhu and M. Wang. 2017. Dissipation, occurrence and risk assessment of a phenylurea herbicide tebuthiuron in sugarcane and aquatic ecosystems in South China. Environ. Pollut. 227: 389–396. https://doi. org/10.1016/j.envpol.2017.04.082
- Roditakis, E., N. Fytrou, M. Staurakaki, J. Vontas and A. Tsagkarakou. 2014. Activity of flonicamid on the sweet potato whitely *Bemisia tabaci* (Homoptera: Aleyrodidae) and its natural enemies. Pest Manag. Sci. 70: 1460–1467. http://dx.doi.org/10.1002/ps.3723
- Saber, A.N., F.M. Malhat, H.M.A. Badawy and D.A. Barakat. 2016. Dissipation dynamic, residue distribution and processing factor of hexythiazox in strawberry fruits under open field condition. Food Chem. 196: 1108–1116. https://doi.org/10.1016/j.foodchem.2015.10.052
- Sadhana, V., K. Senguttuvan, M. Murugan and N. Sathiah. 2021. Efficacy of newer insecticides against whitefly, *Bemisia tabaci* (Hemiptera: Aleyrodidae) on cotton. Madras Agric. J. 107: 1–4. https://doi.org/10.29321/MAJ.10.000334
- Singh, S., L.K. Saini, V.H. Solanki, R.V. Kansara, K.D. Gandhi and N. Patel. 2023. Dissipation kinetics and health risk assessment of certain insecticides applied in/on tomato under open field and poly-house conditions. Heliyon 9: e14963. https://doi.org/10.1016/j. heliyon.2023.e14963
- Sparks, T.C. and R. Nauen. 2015. IRAC: Mode of action classification and insecticide resistance management. Pestic. Biochem. Physiol. 121: 122–128. https://doi.org/10.1016/j. pestbp.2014.11.014
- Subirats, X., S. Reinstadler, S. Porras, M. Raggi and E. Kenndler. 2005. Comparison of methanol and acetonitrile as solvents for the separation of sertindole and its major metabolites by capillary zone electrophoresis. Electrophoresis 26: 3315–3324. https://doi.org/ 10.1002/elps.200500056
- Uesugi, R., A. Jouraku, S.S.N. Pattalung, N. Hinomoto, S. Kuwazaki, H. Kanamori, Y. Katayose and S. Sonoda. 2020. Origin, selection, and spread of diamide insecticide resistance allele in field populations of diamondback moth in east and southeast Asia. Pest Manag. Sci. 77: 313–324. https://doi.org/10.1002/ps.6020
- Ullah, F., H. Gul, N. Desneux, F. Said, X. Gao and D. Song. 2020. Fitness costs in chlorfenapyr resistant populations of the chive maggot, *Bradysia odoriphaga*. Ecotoxicology 29: 407–416. https://doi.org/10.1007/s10646-020-02183-7
- Van Asperen, K.A. 1962. A study of housefly esterases by means of a sensitive colorimetric method. J. Insect Physiol. 8: 401–416. https://doi.org/10.1016/0022-1910(62)90074-4
- Wang, K.Y., Q.L. Guo, X.M. Xia, H.Y. Wang and T.X. Liu. 2007. Resistance of *Aphis gossypii* (Homoptera: Aphididae) to selected insecticides on cotton from five cotton production regions in Shandong, China. J. Pestic. Sci. 32: 372–378. https://doi.org/10.1584/jpestics.G06-51
- Wang, Q., P. Wei, M. Cao, Y. Liu, M. Wang, Y. Guo and G. Zhu. 2016. Residual behavior and risk assessment of the mixed formulation of benzene kresoxim-methyl and fluazinam in cucumber field application. Environ. Monitor. Assess. 188: 341. https://doi.org/10.1007/ s10661-016-5345-9
- Wang, S., F. Jin, X. Cao, Y. Shao, J. Wang, Y. She, Y. Qi, C. Zhang, H. Li, M. Jin, J. Wang, H. Shao and L. Zheng. 2018. Residue behaviors and risk assessment of flonicamid and its metabolites in the cabbage field ecosystem. Ecotoxicol. Environ. Saf. 161: 420–429.
- Zhang, A.J., Y.L. Bian and Y.Z. Feng. 2021. Degradation of chlorfenapyr and thiamethoxam residues in asparagus and risk assessment of dietary intake. Agrochemicals 60: 654–658.
- Zhang, T., Y. Xu, X. Zhou, X. Liang, Y. Bai, F. Sun, W. Zhang, N. Wang, X. Pang and Y. Li. 2022. Dissipation kinetics and safety evaluation of flonicamid in four various types of crops. Molecules 27: 8615. https://doi.org/10.3390/molecules27238615